


(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
7 August 2003 (07.08.2003)

PCT

(10) International Publication Number
WO 03/064596 A2

- (51) International Patent Classification⁷: C12N Westray Drive, Westerville, OH 43081 (US). HUANG, Yung-Sheng; 2462 Danvers Court, Upper Arlington, OH 43220 (US).
- (21) International Application Number: PCT/US03/01698
- (22) International Filing Date: 21 January 2003 (21.01.2003) (74) Agents: BECKER, Cheryl, L. et al.; D-377 AP6A-1, 100 Abbott Park Road, Abbott Park, IL 60064-6008 (US).
- (25) Filing Language: English (81) Designated States (*national*): CA, JP, MX, NO.
- (26) Publication Language: English (84) Designated States (*regional*): European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR).
- (30) Priority Data:
10/060,793 30 January 2002 (30.01.2002) US
- (71) Applicant: ABBOTT LABORATORIES [US/US];
D-377 AP6A-1, 100 Abbott Park Road, Abbott Park, IL 60064-6008 (US).
- (72) Inventors: MUKERJI, Pradip; 1069 Arcaro Drive, Columbus, OH 43230 (US). PEREIRA, Suzette, L.; 710
- Published:
— without international search report and to be republished upon receipt of that report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: DESATURASE GENES, ENZYMES ENCODED THEREBY, AND USES THEREOF



(57) Abstract: Disclosed are isolated polynucleotides encoding an omega-3 desaturase and a delta-12 desaturase, the enzymes encoded by the isolated polynucleotides, vectors containing the isolated polynucleotides, transgenic hosts that contain the isolated polynucleotides that express the enzymes encoded thereby, methods for producing the desaturase enzymes, and method of using the enzymes to make polyunsaturated fatty acids. The isolated polynucleotides are derived from a fungus, *Saprolegnia diclina* (ATCC 56851). In particular, omega-3-desaturase may be utilized, for example, in the conversion of arachidonic acid (AA) to eicosapentaenoic acid (EPA). Delta-12 desaturase may be used, for example, in the conversion of oleic acid (OA) to linoleic (LA). EPA or polyunsaturated fatty acids produced therefrom may be added to pharmaceutical compositions, nutritional compositions, animal feeds, as well as other products such as cosmetics.

DESATURASE GENES, ENZYMES ENCODED THEREBY, AND USES THEREOF

FIELD OF THE INVENTION

The present invention is directed to the identification and isolation of novel genes that encode enzymes involved in the synthesis of polyunsaturated fatty acids (PUFAs). The invention is also directed to the novel desaturase enzymes encoded by these genes and to methods of utilizing the genes and/or the enzymes encoded by the genes. In particular, the invention is directed to genes derived from the fungus *Saprolegnia diclina* (ATCC 56851) that encode a novel ω 3-desaturase (also referred to herein as a Δ 17-desaturase) and a novel Δ 12-desaturase. These enzymes catalyze the introduction of a carbon-carbon double bond between a particular position within a fatty acid substrate. For example, the novel ω 3-desaturase disclosed herein catalyzes the conversion of arachidonic acid (20:4n-6) to eicosapentaenoic acid (20:5n-3) (as well as other desaturation reactions involving other substrates). Likewise, the novel Δ 12-desaturase disclosed herein catalyzes the conversion of oleic acid (18:1n-9) to linoleic acid (18:2n-6). The PUFAs so formed may be added to pharmaceutical compositions, nutritional compositions, animal feeds, or other products.

BACKGROUND

Desaturases are a class of enzymes critical in the production of long-chain polyunsaturated fatty acids. Polyunsaturated fatty acids (PUFAs) play many roles in the proper functioning of all life forms. For example, PUFAs are important components of the plasma membrane of a cell, where they are found in the form of phospholipids. PUFAs also are precursors to mammalian prostacyclins, eicosanoids, leukotrienes and prostaglandins. Additionally, PUFAs are necessary for the proper development of the infant brain, as well as for tissue formation and repair in mature mammals. In view of the biological

significance of PUFAs, attempts are being made to produce them in an efficient manner.

A number of enzymes, most notably desaturases and elongases, are involved in PUFA biosynthesis (see Figure 1). Elongases catalyze the addition of a 2-carbon unit to a fatty acid substrate. Thus, for example, an elongase (generically designated "elo" in Figure 1) catalyzes the conversion of γ -linolenic acid (18:3n-6) to dihomo- γ -linolenic acid (20:3n-6), as well as the conversion of stearidonic acid (18:4n-3) to eicosatetraenoic acid (20:4n-3), etc.

Desaturases catalyze the introduction of unsaturations (i.e., double bonds) between carbon atoms within the fatty acid alkyl chain of the substrate. Thus, for example, linoleic acid (18:2n-6) is produced from oleic acid (18:1n-9) by the action of a Δ 12-desaturase. Similarly, γ -linolenic acid (18:3n-6) is produced from linoleic acid by the action of a Δ 6-desaturase.

Throughout the present application, PUFAs will be unambiguously identified using the "omega" system of nomenclature favored by physiologists and biochemists, as opposed to the "delta" system or I.U.P.A.C. system normally favored by chemists. In the "omega" system, a PUFA is identified by a numeric designation of the number of carbons in the chain. This is followed by a colon and then another numeric designation of the number of unsaturations in the molecule. This is then followed by the designation "n-x," where x is the number of carbons from the methyl end of the molecule where the first unsaturation is located. Each subsequent unsaturation (where there is more than one double bond) is located 3 additional carbon atoms toward the carboxyl end of the molecule. Thus, the PUFAs described herein can be described as being "methylene-interrupted" PUFAs. Where some other designation is required, deviations from the "omega" system will be noted.

Where appropriate, the action of the desaturase enzymes described herein will also be identified using the "omega" system. Thus, an "omega-3" desaturase catalyzes the

introduction of a double bond between the two carbons at positions 3 and 4 from the methyl end of the substrate. However, in many instances, it is more convenient to indicate the activity of a desaturase by counting from the carboxyl end of the substrate. Thus, as shown in Figure 1, a $\Delta 9$ -desaturase catalyzes the introduction of a double bond between the two carbons at positions 9 and 10 from the carboxyl end of the substrate. In short, where the term " ω " is used, the position on the substrate is being designated relative to the methyl terminus; where the term " Δ " is used, the position on the substrate is being designated relative to the carboxyl terminus.

It must be noted that mammals cannot desaturate fatty acid substrates beyond the $\Delta 9$ position (i.e., beyond 9 carbon atoms distant from the carboxyl terminus). Thus, for example, mammals cannot convert oleic acid (18:1n-9) into linoleic acid (18:2n-6); linoleic acid contains an unsaturation at position $\Delta 12$. Likewise, α -linolenic acid (18:3n-3) (having unsaturations at $\Delta 12$ and $\Delta 15$) cannot be synthesized by mammals. However, for example, mammals can convert α -linolenic acid into stearidonic acid (18:4n-3) by the action of a $\Delta 6$ -desaturase. (See Figure 1. See also PCT publication WO 96/13591; The FASEB Journal, Abstracts, Part I, Abstract 3093, page A532 (Experimental Biology 98, San Francisco, CA, April 18-22, 1998); and U.S. Patent No. 5,552,306.)

Still referring to Figure 1, in mammals, fungi, and algae, the stearidonic acid so formed is converted into eicosatetraenoic acid (20:4n-3) by the action of an elongase. This PUFA can then be converted to eicosapentaenoic acid (20:5n-3) by a $\Delta 5$ -desaturase. Eicosapentaenoic acid can then, in turn, be converted to $\omega 3$ -docosapentaenoic acid (22:5n-3) by an elongase.

Other eukaryotes, including fungi and plants, have enzymes that desaturate fatty acid substrates at carbon $\Delta 12$ (see PCT

publication WO 94/11516 and U.S. Patent No. 5,443,974) and at carbon delta-15 (see PCT publication WO 93/11245). The major polyunsaturated fatty acids of animals therefore are either derived from diet and/or from desaturation and elongation of linoleic acid or α -linolenic acid. In view of these difficulties, there remains a significant need to isolate genes involved in PUFA synthesis. Ideally, these genes would originate from species that naturally produce fatty acids that are not produced naturally in mammals. These genes could then be expressed in a microbial, plant, or animal system, which would thereby be altered to produce commercial quantities of one or more PUFAs. Thus, there is a definite need for novel $\Delta 12$ - and $\Delta 17$ -desaturase enzymes, the respective genes encoding these enzymes, as well as recombinant methods of producing these enzymes. Additionally, a need exists for oils containing levels of PUFAs beyond those naturally present. Access to such $\Delta 12$ - and $\Delta 17$ -desaturase enzymes allows for the production of large amounts of PUFAs that cannot be synthesized *de novo* in mammals. These PUFAs can be used as pharmaceutical agents and/or nutritional supplements.

All patents, patent publications and priority documents cited herein are hereby incorporated by reference in their entirety.

SUMMARY OF THE INVENTION

One embodiment of the present invention encompasses an isolated nucleotide acid sequence or fragment thereof comprising or complementary to a nucleotide sequence encoding a polypeptide having desaturase activity, wherein the amino acid sequence of the polypeptide has at least 50% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:26 and SEQ ID NO:42.

The present invention also includes an isolated nucleotide

sequence (or fragment thereof) comprising or complementary to at least 50% of the nucleotide sequence selected from the group consisting of SEQ ID NO:25 and SEQ ID NO:41. In particular, the sequence may be selected from the group consisting of SEQ ID NO:25 and SEQ ID NO:41. The sequence may encode a functionally active desaturase which utilizes a polyunsaturated fatty acid as a substrate. Furthermore, the nucleotide sequence may be isolated from a fungus, such as *Saprolegnia diclina*.

An additional embodiment of the present invention includes a purified polypeptide encoded by the nucleotide sequences described above.

The present invention also includes a purified polypeptide that desaturates a polyunsaturated fatty acid substrate at an omega-3 carbon of the substrate and has at least 50% amino acid identity to an amino acid sequence comprising SEQ ID NO: 26. The Polypeptide may desaturate a fatty acid substrate having 20 carbon atoms.

Additionally, the present invention encompasses a purified polypeptide that desaturates a polyunsaturated fatty acid substrate at a delta-12 carbon of the substrate and has at least 50% amino acid identity to SEQ ID NO: 42. The polypeptide may desaturate a fatty acid substrate having 18 carbon atoms.

Another embodiment of the present invention includes a method of producing a desaturase comprising the steps of: isolating a nucleotide sequence comprising or complementary to at least 50% of the nucleotide sequence selected from the group consisting of SEQ ID NO: 25 and SEQ ID NO: 41; constructing a vector comprising the isolated nucleotide sequence; and introducing the vector into a host cell for a time and under conditions sufficient for expression of a desaturase encoded by the isolated nucleotide sequence.

A further embodiment of the present invention includes a vector comprising: 1) an isolated nucleotide sequence corresponding to or complementary to at least about 50% of the nucleotide sequence selected from the group consisting of SEQ ID

NO: 25 and SEQ ID NO: 41, operably linked to b) a regulatory sequence.

Additionally, another embodiment of the present invention includes a host cell comprising the above vector. The host cell may be, for example, a eukaryotic cell selected from the group consisting of a mammalian cell, an insect cell, a plant cell and a fungal cell. With respect to the host cell, expression of the isolated nucleotide sequence of the vector may result in the host cell producing a polyunsaturated fatty acid that is not produced in a wild-type of the host cell.

Also, the present invention encompasses a plant cell, plant, or plant tissue comprising the vector described above, wherein expression of the nucleotide sequence of the vector results in production of a polyunsaturated fatty acid by the plant cell, plant or plant tissue. The vector in the plant cell, plant or plant tissue may induce the production of a polyunsaturated fatty acid selected from the group consisting of, for example, linoleic acid, eicosatetraenoic acid and eicosapentaenoic acid. Also, the invention includes one or more plant oils or acids expressed by the plant cell, plant or plant tissue.

The invention also includes a transgenic plant comprising the above-described vector, wherein expression of the nucleotide sequence of the vector results in production of a polyunsaturated fatty acid in seeds of the transgenic plant.

Another embodiment of the present invention encompasses a method for producing a polyunsaturated fatty acid comprising the steps of: isolating a nucleotide sequence comprising or complementary to at least about 50% of the nucleotide sequence selected from the group consisting of SEQ ID NO: 25 and SEQ ID NO: 41; constructing a vector comprising the isolated nucleotide sequence; transforming the vector into a host cell for a time and under conditions sufficient for expression of a desaturase encoded by the isolated nucleotide sequence; and exposing the expressed desaturase selected from the group consisting of an omega-3-desaturase and a delta 12-desaturase, to a fatty acid substrate, whereby the substrate is catalytically converted by

said desaturase into a desired polyunsaturated fatty acid product. The substrate is dihomogamma-linolenic acid or arachidonic acid and the product polyunsaturated fatty acid is eicosatetraenoic acid or eicosapentaenoic acid, respectively, when the expressed desaturase is an omega-3-desaturase. The substrate polyunsaturated fatty acid is oleic acid and the product polyunsaturated fatty acid is linoleic acid, when the expressed desaturase is a delta 12-desaturase.

The method may further comprise the step of exposing the polyunsaturated fatty acid product to one or more enzymes selected from the group consisting of a desaturase and an elongase, whereby the polyunsaturated fatty acid product is catalytically converted into another polyunsaturated fatty acid product. The product polyunsaturated fatty acid is eicosatetraenoic acid or eicosapentaenoic acid and the another polyunsaturated fatty acid is eicosapentaenoic acid or omega 3-docosapentaenoic acid, respectively, when the expressed desaturase is an omega 3-desaturase. The product polyunsaturated fatty acid is linoleic acid and the another polyunsaturated fatty acid is gamma-linolenic acid, when the expressed desaturase is a delta 12-desaturase.

Additionally, the method described directly above may further comprise the step of exposing the another polyunsaturated fatty acid to one or more enzymes selected from the group consisting of a desaturase and an elongase in order to convert the another polyunsaturated fatty acid to a final polyunsaturated fatty acid. The final polyunsaturated fatty acid is selected from the group consisting of omega 3-docosapentaenoic acid and docosahexaenoic acid, when the expressed desaturase of step (d) is an omega 3-desaturase. In contrast, the final polyunsaturated fatty acid is selected from the group consisting of dihomogamma-linolenic acid, arachidonic acid, adrenic acid, omega 6-docosapentaenoic acid, eicosatetraenoic acid, stearidonic acid, eicosapentaenoic acid, omega 3-docosapentaenoic acid and docosahexaenoic acid, when the expressed desaturase is a delta 12-desaturase.

An additional embodiment of the present invention includes a method of producing a polyunsaturated fatty acid comprising exposing a fatty acid substrate to a polypeptide having at least 50% amino acid identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 26 and SEQ ID NO: 42, whereby the fatty acid substrate is catalytically converted into the polyunsaturated fatty acid product. The fatty acid substrate is dihomo-gamma-linolenic acid or arachidonic acid and the product polyunsaturated fatty acid is eicosatetraenoic acid or eicosapentaenoic acid, respectively, when the polypeptide is an omega 3-desaturase. In contrast, the fatty acid substrate is oleic acid and the polyunsaturated fatty acid product is linoleic acid, when the polypeptide is a delta 12-desaturase.

A further embodiment of the present invention includes a composition comprising at least one polyunsaturated fatty acid selected from the group consisting of the "product" polyunsaturated fatty acid produced according to the method described above, the "another" polyunsaturated fatty acid produced according to the method described above, and the "final" polyunsaturated fatty acid produced according to the method described above. The product polyunsaturated fatty acid is eicosatetraenoic acid or eicosapentaenoic acid, when the expressed desaturase is an omega 3-desaturase. In contrast, the product polyunsaturated fatty acid is linoleic acid, when the expressed desaturase is a delta 12-desaturase. The another polyunsaturated fatty acid is eicosapentaenoic acid or omega 3-docosapentaenoic acid, respectively, when the expressed desaturase is an omega 3-desaturase. However, the another polyunsaturated fatty acid is gamma-linolenic acid, when the expressed desaturase is a delta 12-desaturase. The final polyunsaturated fatty acid is selected from the group consisting of omega 3-docosapentaenoic acid and docosahexaenoic acid, when the expressed desaturase is an omega 3-desaturase. In contrast, the final polyunsaturated fatty acid is selected from the group consisting of dihomo-gamma-linolenic acid, arachidonic acid, adrenic acid, omega 6-docosapentaenoic acid, eicosatetraenoic

acid, stearidonic acid, eicosapentaenoic acid, omega 3-docosapentaenoic acid and docosahexaenoic acid, when the expressed desaturase is a delta 12-desaturase.

A further embodiment of the present invention includes a method of preventing or treating a condition caused by insufficient intake of at least one polyunsaturated fatty acid comprising administering to the patient the above-described composition in an amount sufficient to effect the prevention or treatment.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic showing the biosynthetic pathway leading to the production of various PUFAs.

Figure 2 is the nucleotide sequence of *sdd17* (SEQ ID NO: 25), a gene derived from *S. diclina* (ATCC 56851) that encodes a novel ω 3-fatty acid desaturase.

Figure 3 is the amino acid sequence of the ω 3-desaturase (SDD17) (SEQ ID NO: 26) encoded by the nucleotide sequence depicted in Figure 2.

Figure 4 is an amino acid sequence comparison between the SDD17 desaturase depicted in Figure 3 and a known Δ 15-desaturase from *Synechocystis* sp. (SYCDESB).

Figure 5 is an amino acid sequence comparison between the SDD17 desaturase depicted in Figure 3 and a known Δ 17-desaturase from *C. elegans* (CELEFAT).

Figure 6 is the nucleotide sequence of *sdd12* (SEQ ID NO:41), a gene derived from *S. diclina* (ATCC 56851) that encodes a novel Δ 12-fatty acid desaturase.

Figure 7 is the amino acid sequence of the Δ 12-desaturase (SDD12) (SEQ ID NO: 42) encoded by the nucleotide sequence depicted in Figure 6.

Figure 8 is an amino acid sequence comparison between the SDD12 desaturase depicted in Figure 7 and a known $\Delta 12$ -desaturase from *G. hirsutum* (GHO6DES).

Figure 9 lists the sequence identifiers used throughout the application as well as the corresponding amino acid or nucleotide sequence.

DETAILED DESCRIPTION OF THE INVENTION

Abbreviations utilized herein:

- 18:1n-9 = oleic acid = OA
- 18:2n-6 = linoleic acid = LA
- 18:3n-6 = gamma-linolenic acid = GLA
- 18:3n-3 = alpha-linolenic acid = ALA
- 18:4n-3 = stearidonic acid = STA
- 20:3n-6 = dihomo-gamma-linolenic acid = DGLA
- 20:4n-6 = arachidonic acid = AA
- 20:4n-3 = eicosatetraenoic acid = ETA
- 20:5n-3 = eicosapentaenoic acid = EPA
- 22:4n-6 = adrenic acid
- 22:5n-3 = omega-3-docosapentaenoic acid = DPA
- 22:6n-3 = docosahexaenoic acid = DHA
- PUFA = polyunsaturated fatty acid

The subject invention relates to the nucleotide and translated amino acid sequences of the $\omega 3$ -desaturase and $\Delta 12$ -desaturase genes isolated from the fungus *Saprolegnia diclina* or *S. diclina* (ATCC 56851). Furthermore, the subject invention also includes uses of these genes and of the enzymes encoded by these genes. For example, the genes and their corresponding enzymes may be used in the production of polyunsaturated fatty acids such as linoleic acid, eicosapentaenoic acid, and the like. These fatty acids can be added to pharmaceutical compositions, nutritional compositions, and to other valuable products.

The fungus *S. diclina* (ATCC 56851), from which the polynucleotides described herein were isolated, is available commercially from the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110. The fungus is supplied frozen and can be propagated in ATCC medium 307 cornmeal agar (Difco # 0386) at 24°C. For further information on this fungus, see Beakes G. (1983) "A comparative account of cyst coat ontogeny in saprophytic and fish-lesion (pathogenic) isolates of the *Saprolegnia diclina*-*parasitica* complex." *Can. J. Bot.* 61, 603-625; and Willoughby L.G., et al. (1983) "Zoospore germination of *Saprolegnia* pathogenic to fish." *Trans. Br. Mycol. Soc.* 80, 421-435.

The ω 3-Desaturase Gene, the Δ 12-Desaturase Gene, and the Enzymes Encoded Thereby

The enzymes encoded by the omega-3 desaturase and delta-12 desaturase genes of the present invention are essential in the production of PUFAs having at least two unsaturations and an overall length of 18 carbons or longer. The nucleotide sequence of the isolated *Saprolegnia diclina* omega-3 desaturase gene is shown in SEQ ID NO: 25 and in Figure 2. This gene differs significantly in sequence from all known desaturase genes, from any source. The encoded omega-3 desaturase enzyme is shown in SEQ ID NO: 26 and in Figure 3. The nucleotide sequence of the isolated *Saprolegnia diclina* delta-12 desaturase gene is shown in SEQ ID NO: 41 and in Figure 6. This gene also differs significantly in sequence from all known desaturase genes, from any source. The encoded delta-12 desaturase enzyme is shown in SEQ ID NO: 42 and in Figure 7.

The isolated omega-3 desaturase gene of the present invention, when transformed into a yeast host, produces an omega-3 desaturase enzyme that readily catalyzes the conversion of DGLA to ETA, AA to EPA, and adrenic acid to DPA (see Example 5). In like manner, the isolated delta-12 desaturase gene of the present

invention, when transformed into a yeast host, produces a delta-12 desaturase enzyme that readily catalyzes the conversion OA to LA (see Example 9).

It should be noted that the present invention also encompasses nucleotide sequences (and the corresponding encoded proteins) having sequences comprising, identical to, or complementary to at least about 50%, preferably at least about 60%, and more preferably at least about 70%, even more preferably at least about 80%, and most preferably at least about 90% of the nucleotides (i.e., having sequence identity to the sequence) shown in SEQ ID NO: 25 and SEQ ID NO: 41 (i.e., the nucleotide sequences of the omega-3 desaturase gene and the delta-12 desaturase gene of *Saprolegnia diclina*, respectively) described herein. (All integers between 50% and 100% are also considered to be within the scope of the present invention with respect to percent identity.) Such sequences may be derived from any source, either isolated from a natural source, or produced via a semi-synthetic route, or synthesized *de novo*. Such sequences may be isolated from or derived from fungal sources, as well as other non-fungal sources, such as bacterial, algal, *C. elegans*, mouse or human.

The present invention also encompasses fragments and derivatives of the nucleotide sequences shown in SEQ ID NO: 25 and SEQ ID NO: 41, as well as fragments and derivatives of the sequences derived from other sources, and having the above-described complementarity, identity or correspondence. Functional equivalents of the above-sequences (i.e., sequences having omega-3 desaturase activity or delta-12 desaturase activity, as appropriate) are also encompassed by the present invention.

For purposes of the present invention, a "fragment" of a nucleotide sequence is defined as a contiguous sequence of approximately at least 6, preferably at least about 8, more preferably at least about 10 nucleotides, and even more

preferably at least about 15 nucleotides corresponding to a region of the specified nucleotide sequence.

Sequence identity or percent identity is the number of exact matches between two aligned sequences divided by the length of the shorter sequence and multiplied by 100. An approximate alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2:482-489 (1981). This algorithm may be extended to use with peptide or protein sequences using the scoring matrix created by Dayhoff, *Atlas of Protein Sequences and Structure*, M.O. Dayhoff ed., 5 Suppl. 3:353-358, National Biomedical Research Foundation, Washington, D.C., USA, and normalized by Gribskov, *Nucl. Acids Res.* 14(6):6745-66763 (1986). The Genetics Computer Group (GCG) (Madison, Wisconsin) provides a computer program that automates this algorithm for both nucleic acid and peptide sequences in the "BestFit" utility application. The default parameters for this method are described in the Wisconsin Sequence Analysis Package Program Manual, Version 8 (1995) (available from GCG). Other equally suitable programs for calculating the percent identity or similarity between sequences are generally known in the art.

The invention also includes a purified polypeptide which desaturates PUFAs at the omega-3 position and has at least about 50% amino acid similarity or identity, preferably at least about 60% similarity or identity, more preferably at least about 70% similarity or identity, even more preferably at least about 80% similarity or identity, and most preferably at least about 90% similarity or identity to the amino acid sequence shown in SEQ ID NO: 26 (Figure 3) and encoded by the above-noted nucleotide sequence(s) (All integers between 50% and 100% similarity or identity are also included within the scope of the invention.)

The invention further includes a purified polypeptide which desaturates PUFAs at the delta-12 position and has at least about 50% amino acid similarity or identity, preferably at least about 60% similarity or identity, more preferably at least about 70% similarity or identity, even more preferably at least about 80%

similarity or identity, and most preferably at least about 90% similarity or identity to the amino acid sequence shown in SEQ. ID. NO: 42 (Figure 7) which, in turn, is encoded by the above-described nucleotide sequence(s). (All integers between 50% and 100% similarity or identity are also included within the scope of the invention.)

The term "identity" refers to the relatedness of two sequences on a nucleotide-by-nucleotide basis over a particular comparison window or segment. Thus, identity is defined as the degree of sameness, correspondence or equivalence between the same strands (either sense or antisense) of two DNA segments. "Percentage of sequence identity" is calculated by comparing two optimally aligned sequences over a particular region, determining the number of positions at which the identical base occurs in both sequence in order to yield the number of matched positions, dividing the number of such positions by the total number of positions in the segment being compared and multiplying the result by 100. Optimal alignment of sequences may be conducted by the algorithm of Smith & Waterman, Appl. Math. 2:482 (1981), by the algorithm of Needleman & Wunsch, J. Mol. Biol. 48:443 (1970), by the method of Pearson & Lipman, Proc. Natl. Acad. Sci. (USA) 85:2444 (1988) and by computer programs which implement the relevant algorithms (e.g., Clustal Macaw Pileup (<http://cmgm.stanford.edu/biochem218/11Multiple.pdf>; Higgins et al., CABIOS. 5L151-153 (1989)), FASTDB (Intelligenetics), BLAST (National Center for Biomedical Information; Altschul et al., Nucleic Acids Research 25:3389-3402 (1997)), PILEUP (Genetics Computer Group, Madison, WI) or GAP, BESTFIT, FASTA and TFASTA (Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, Madison, WI). (See U.S. Patent No. 5,912,120.)

For purposes of the present invention, "complementarity" is defined as the degree of relatedness between two DNA segments. It is determined by measuring the ability of the sense strand of one DNA segment to hybridize with the antisense strand of the other DNA segment, under appropriate conditions, to form a double helix. A "complement" is defined as a sequence which pairs to a given sequence based upon the canonic base-pairing rules. For example, a sequence A-G-T in one nucleotide strand is "complementary" to T-C-A in the other strand.

In the double helix, adenine appears in one strand, thymine appears in the other strand. Similarly, wherever guanine is found in one strand, cytosine is found in the other. The greater the relatedness between the nucleotide sequences of two DNA segments, the greater the ability to form hybrid duplexes between the strands of the two DNA segments.

"Similarity" between two amino acid sequences is defined as the presence of a series of identical as well as conserved amino acid residues in both sequences. The higher the degree of similarity between two amino acid sequences, the higher the correspondence, sameness or equivalence of the two sequences. ("Identity between two amino acid sequences is defined as the presence of a series of exactly alike or invariant amino acid residues in both sequences.") The definitions of "complementarity", "identity" and "similarity" are well known to those of ordinary skill in the art.

"Encoded by" refers to a nucleic acid sequence which codes for a polypeptide sequence, wherein the polypeptide sequence or a portion thereof contains an amino acid sequence of at least 3 amino acids, more preferably at least 8 amino acids, and even more preferably at least 15 amino acids from a polypeptide encoded by the nucleic acid sequence.

The present invention also encompasses an isolated nucleotide sequence which encodes PUFA desaturase activity and that is hybridizable, under moderately stringent conditions, to a nucleic acid having a nucleotide sequence comprising or complementary to the nucleotide sequence comprising SEQ ID NO:25 or SEQ ID NO:41. A nucleic acid molecule is "hybridizable" to another nucleic acid molecule when a single-stranded form of the nucleic acid molecule can anneal to the other nucleic acid molecule under the appropriate conditions of temperature and ionic strength (see Sambrook et al., "Molecular Cloning: A Laboratory Manual, Second Edition (1989), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York)). The conditions of temperature and ionic strength determine the "stringency" of the hybridization. "Hybridization" requires that two nucleic acids contain complementary sequences. However, depending on the stringency of the hybridization, mismatches between bases may occur. The appropriate stringency for hybridizing nucleic acids depends on the length of the nucleic acids and the degree of complementation. Such variables are well known in the art. More specifically, the greater the degree of similarity or homology between two nucleotide sequences, the greater the value of T_m for hybrids of nucleic acids having those sequences. For hybrids of greater than 100 nucleotides in length, equations for calculating T_m have been derived (see Sambrook et al., *supra*). For hybridization with shorter nucleic acids, the position of mismatches becomes more important, and the length of the oligonucleotide determines its specificity (see Sambrook et al., *supra*).

As used herein, an "isolated nucleic acid fragment or sequence" is a polymer of RNA or DNA that is single- or double-stranded, optionally containing synthetic, non-natural or altered nucleotide bases. An isolated nucleic acid fragment in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA or synthetic DNA. (A "fragment" of a specified polynucleotide refers to a polynucleotide sequence

which comprises a contiguous sequence of approximately at least about 6 nucleotides, preferably at least about 8 nucleotides, more preferably at least about 10 nucleotides, and even more preferably at least about 15 nucleotides, and most preferable at least about 25 nucleotides identical or complementary to a region of the specified nucleotide sequence.) Nucleotides (usually found in their 5'-monophosphate form) are referred to by their single letter designation as follows: "A" for adenylate or deoxyadenylate (for RNA or DNA, respectively), "C" for cytidylate or deoxycytidylate, "G" for guanylate or deoxyguanylate, "U" for uridylate, "T" for deoxythymidylate, "R" for purines (A or G), "Y" for pyrimidines (C or T), "K" for G or T, "H" for A or C or T, "I" for inosine, and "N" for any nucleotide.

The terms "fragment or subfragment that is functionally equivalent" and "functionally equivalent fragment or subfragment" are used interchangeably herein. These terms refer to a portion or subsequence of an isolated nucleic acid fragment in which the ability to alter gene expression or produce a certain phenotype is retained whether or not the fragment or subfragment encodes an active enzyme. For example, the fragment or subfragment can be used in the design of chimeric constructs to produce the desired phenotype in a transformed plant. Chimeric constructs can be designed for use in co-suppression or antisense by linking a nucleic acid fragment or subfragment thereof, whether or not it encodes an active enzyme, in the appropriate orientation relative to a plant promoter sequence.

The terms "homology", "homologous", "substantially similar" and "corresponding substantially" are used interchangeably herein. They refer to nucleic acid fragments wherein changes in one or more nucleotide bases does not affect the ability of the nucleic acid fragment to mediate gene expression or produce a certain phenotype. These terms also refer to modifications of the nucleic acid fragments of the instant invention such as deletion or insertion of one or more nucleotides that do not substantially alter the functional

properties of the resulting nucleic acid fragment relative to the initial, unmodified fragment. It is therefore understood, as those skilled in the art will appreciate, that the invention encompasses more than the specific exemplary sequences.

``Gene'' refers to a nucleic acid fragment that expresses a specific protein, including regulatory sequences preceding (5' non-coding sequences) and following (3' non-coding sequences) the coding sequence.

``Native gene'' refers to a gene as found in nature with its own regulatory sequences. In contrast, ``chimeric construct'' refers to a combination of nucleic acid fragments that are not normally found together in nature. Accordingly, a chimeric construct may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner different than that normally found in nature. (The term ``isolated'' means that the sequence is removed from its natural environment.)

A ``foreign'' gene refers to a gene not normally found in the host organism, but that is introduced into the host organism by gene transfer. Foreign genes can comprise native genes inserted into a non-native organism, or chimeric constructs. A ``transgene'' is a gene that has been introduced into the genome by a transformation procedure.

``Coding sequence'' refers to a DNA sequence that codes for a specific amino acid sequence. ``Regulatory sequences'' refer to nucleotide sequences located upstream (5' non-coding sequences), within, or downstream (3' non-coding sequences) of a coding sequence, and which influence the transcription, RNA processing or stability, or translation of the associated coding sequence. Regulatory sequences may include, but are not limited to, promoters, translation leader sequences, introns, and polyadenylation recognition sequences.

``Promoter'' refers to a DNA sequence capable of controlling the expression of a coding sequence or functional RNA. The promoter sequence consists of proximal and more distal upstream

elements, the latter elements often referred to as enhancers. Accordingly, an "enhancer" is a DNA sequence which can stimulate promoter activity and may be an innate element of the promoter or a heterologous element inserted to enhance the level or tissue-specificity of a promoter. Promoter sequences can also be located within the transcribed portions of genes, and/or downstream of the transcribed sequences. Promoters may be derived in their entirety from a native gene, or be composed of different elements derived from different promoters found in nature, or even comprise synthetic DNA segments. It is understood by those skilled in the art that different promoters may direct the expression of a gene in different tissues or cell types, or at different stages of development, or in response to different environmental conditions. Promoters which cause a gene to be expressed in most cell types at most times are commonly referred to as "constitutive promoters". New promoters of various types useful in plant cells are constantly being discovered; numerous examples may be found in the compilation by Okamuro and Goldberg, (1989) *Biochemistry of Plants* 15:1-82. It is further recognized that since in most cases the exact boundaries of regulatory sequences have not been completely defined, DNA fragments of some variation may have identical promoter activity.

An "intron" is an intervening sequence in a gene that does not encode a portion of the protein sequence. Thus, such sequences are transcribed into RNA but are then excised and are not translated. The term is also used for the excised RNA sequences. An "exon" is a portion of the sequence of a gene that is transcribed and is found in the mature messenger RNA derived from the gene, but is not necessarily a part of the sequence that encodes the final gene product.

The "translation leader sequence" refers to a DNA sequence located between the promoter sequence of a gene and the coding sequence. The translation leader sequence is present in the fully processed mRNA upstream of the translation start sequence. The translation leader sequence may affect processing of the

primary transcript to mRNA, mRNA stability or translation efficiency. Examples of translation leader sequences have been described (Turner, R. and Foster, G. D. (1995) *Molecular Biotechnology* 3:225).

The "3' non-coding sequences" refer to DNA sequences located downstream of a coding sequence and include polyadenylation recognition sequences and other sequences encoding regulatory signals capable of affecting mRNA processing or gene expression. The polyadenylation signal is usually characterized by affecting the addition of polyadenylic acid tracts to the 3' end of the mRNA precursor. The use of different 3' non-coding sequences is exemplified by Ingelbrecht et al., (1989) *Plant Cell* 1:671-680.

"RNA transcript" refers to the product resulting from RNA polymerase-catalyzed transcription of a DNA sequence. When the RNA transcript is a perfect complementary copy of the DNA sequence, it is referred to as the primary transcript or it may be a RNA sequence derived from post-transcriptional processing of the primary transcript and is referred to as the mature RNA.

"Messenger RNA (mRNA)" refers to the RNA that is without introns and that can be translated into protein by the cell.

"cDNA" refers to a DNA that is complementary to and synthesized from a mRNA template using the enzyme reverse transcriptase.

The cDNA can be single-stranded or converted into the double-stranded form using the Klenow fragment of DNA polymerase I.

"Sense" RNA refers to RNA transcript that includes the mRNA and can be translated into protein within a cell or *in vitro*.

"Antisense RNA" refers to an RNA transcript that is complementary to all or part of a target primary transcript or mRNA and that blocks the expression of a target gene (U.S. Patent No. 5,107,065). The complementarity of an antisense RNA may be with any part of the specific gene transcript, i.e., at the 5' non-coding sequence, 3' non-coding sequence, introns, or the coding sequence. "Functional RNA" refers to antisense RNA, ribozyme RNA, or other RNA that may not be translated but yet has an effect on cellular processes. The terms "complement"

and "reverse complement" are used interchangeably herein with respect to mRNA transcripts, and are meant to define the antisense RNA of the message.

The term "endogenous RNA" refers to any RNA which is encoded by any nucleic acid sequence present in the genome of the host prior to transformation with the recombinant construct of the present invention, whether naturally-occurring or non-naturally occurring, i.e., introduced by recombinant means, mutagenesis, etc.

The term "non-naturally occurring" means artificial, not consistent with what is normally found in nature.

The term "operably linked" refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is regulated by the other. For example, a promoter is operably linked with a coding sequence when it is capable of regulating the expression of that coding sequence (i.e., that the coding sequence is under the transcriptional control of the promoter). Coding sequences can be operably linked to regulatory sequences in a sense or antisense orientation. In another example, the complementary RNA regions of the invention can be operably linked, either directly or indirectly, 5' to the target mRNA, or 3' to the target mRNA, or within the target mRNA, or a first complementary region is 5' and its complement is 3' to the target mRNA.

The term "expression", as used herein, refers to the production of a functional end-product. Expression of a gene involves transcription of the gene and translation of the mRNA into a precursor or mature protein. "Antisense inhibition" refers to the production of antisense RNA transcripts capable of suppressing the expression of the target protein.

"Co-suppression" refers to the production of sense RNA transcripts capable of suppressing the expression of identical or substantially similar foreign or endogenous genes (U.S. Patent No. 5,231,020).

"Mature" protein refers to a post-translationally processed polypeptide; i.e., one from which any pre- or

propeptides present in the primary translation product have been removed. "Precursor" protein refers to the primary product of translation of mRNA; i.e., with pre- and propeptides still present. Pre- and propeptides may be but are not limited to intracellular localization signals.

"Stable transformation" refers to the transfer of a nucleic acid fragment into a genome of a host organism, including both nuclear and organellar genomes, resulting in genetically stable inheritance. In contrast, "transient transformation" refers to the transfer of a nucleic acid fragment into the nucleus, or DNA-containing organelle, of a host organism resulting in gene expression without integration or stable inheritance. Host organisms containing the transformed nucleic acid fragments are referred to as "transgenic" organisms. The preferred method of cell transformation of rice, corn and other monocots is the use of particle-accelerated or "gene gun" transformation technology (Klein et al., (1987) *Nature (London)* 327:70-73; U.S. Patent No. 4,945,050), or an *Agrobacterium*-mediated method using an appropriate Ti plasmid containing the transgene (Ishida Y. et al., 1996, *Nature Biotech.* 14:745-750). The term "transformation" as used herein refers to both stable transformation and transient transformation.

Standard recombinant DNA and molecular cloning techniques used herein are well known in the art and are described more fully in Sambrook, J., Fritsch, E.F. and Maniatis, T. *Molecular Cloning: A Laboratory Manual*; Cold Spring Harbor Laboratory Press: Cold Spring Harbor, 1989 (hereinafter "Sambrook").

The term "recombinant" refers to an artificial combination of two otherwise separated segments of sequence, e.g., by chemical synthesis or by the manipulation of isolated segments of nucleic acids by genetic engineering techniques.

"PCR" or "Polymerase Chain Reaction" is a technique for the synthesis of large quantities of specific DNA segments, consists of a series of repetitive cycles (Perkin Elmer Cetus Instruments, Norwalk, CT). Typically, the double stranded DNA is heat denatured, the two primers complementary to the

3' boundaries of the target segment are annealed at low temperature and then extended at an intermediate temperature. One set of these three consecutive steps is referred to as a cycle.

Polymerase chain reaction ("PCR") is a powerful technique used to amplify DNA millions of fold, by repeated replication of a template, in a short period of time. (Mullis et al, *Cold Spring Harbor Symp. Quant. Biol.* 51:263-273 (1986); Erlich et al, European Patent Application 50,424; European Patent Application 84,796; European Patent Application 258,017, European Patent Application 237,362; Mullis, European Patent Application 201,184, Mullis et al U.S. Patent No. 4,683,202; Erlich, U.S. Patent No. 4,582,788; and Saiki et al, U.S. Patent No. 4,683,194). The process utilizes sets of specific in vitro synthesized oligonucleotides to prime DNA synthesis. The design of the primers is dependent upon the sequences of DNA that are desired to be analyzed. The technique is carried out through many cycles (usually 20-50) of melting the template at high temperature, allowing the primers to anneal to complementary sequences within the template and then replicating the template with DNA polymerase.

The products of PCR reactions are analyzed by separation in agarose gels followed by ethidium bromide staining and visualization with UV transillumination. Alternatively, radioactive dNTPs can be added to the PCR in order to incorporate label into the products. In this case the products of PCR are visualized by exposure of the gel to x-ray film. The added advantage of radiolabeling PCR products is that the levels of individual amplification products can be quantitated.

The terms "recombinant construct", "expression construct" and "recombinant expression construct" are used interchangeably herein. These terms refer to a functional unit of genetic material that can be inserted into the genome of a cell using standard methodology well known to one skilled in the art. Such construct may be itself or may be used in conjunction with a vector. If a vector is used then the choice of vector is

dependent upon the method that will be used to transform host plants as is well known to those skilled in the art. For example, a plasmid vector can be used. The skilled artisan is well aware of the genetic elements that must be present on the vector in order to successfully transform, select and propagate host cells comprising any of the isolated nucleic acid fragments of the invention. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones et al., (1985) *EMBO J.* 4:2411-2418; De Almeida et al., (1989) *Mol. Gen. Genetics* 218:78-86), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, Western analysis of protein expression, or phenotypic analysis.

Expression of the Omega-3-Desaturase and the Delta 12-Desaturase Genes

Once the genes encoding the omega-3 and delta-12 desaturase enzymes have been isolated, they may then be introduced into either a prokaryotic or eukaryotic host cell (individually or in combination) through the use of a vector or construct. The vector, for example, a bacteriophage, cosmid or plasmid, may comprise the nucleotide sequence encoding either or both of the desaturase enzymes, as well as any promoter which is functional in the host cell and is able to elicit expression of the desaturase(s) encoded by the nucleotide sequence(s). The promoter is in operable association with, or operably linked, to the nucleotide sequence. (As noted above, a regulatory sequence (e.g., a promoter) is said to be "operably linked" with a coding sequence if the regulatory sequence affects transcription or expression of the coding sequence. The promoter (or other type of regulatory sequence) need not be directly linked to the coding sequence. Suitable promoters include, for example, those from genes encoding alcohol dehydrogenase, glyceraldehyde-3 phosphate dehydrogenase, phosphoglucosomerase, phosphoglycerate

kinase, acid phosphatase, T7, TPI, lactase, metallothionein, cytomegalovirus immediate early, whey acidic protein, glucoamylase, and promoters activated in the presence of galactose, for example, GAL1 and GAL10. Additionally, nucleotide sequences which encode other proteins, oligosaccharides, lipids, etc. may also be included within the vector as well as other regulatory sequences such as a polyadenylation signal (e.g., the poly-A signal of SV-40T-antigen, ovalalbumin or bovine growth hormone), antibiotic resistance markers, auxotrophic markers, and the like. The choice of sequences present in the construct is dependent upon the desired expression products, the nature of the host cell, and the proposed means to separate transformed cells from non-transformed cells.

As noted above, once the vector has been constructed, it may then be introduced into the host cell of choice by methods known to those of ordinary skill in the art including, for example, transfection, transformation and electroporation (see Molecular Cloning: A Laboratory Manual, 2nd ed., Vol. 1-3, ed. Sambrook et al., Cold Spring Harbor Laboratory Press (1989)). The host cell is then cultured under suitable conditions permitting expression of the genes leading to the production of the desired PUFA, which is then recovered and purified. (The substrates which may be produced by the host cell either naturally or transgenically, as well as the enzymes which may be encoded by DNA sequences present in the vector, which is subsequently introduced into the host cell, are shown in Figure 1.)

Examples of suitable prokaryotic host cells include, for example, bacteria such as *Escherichia coli* and *Bacillus subtilis*, as well as cyanobacteria such as *Spirulina* spp. (i.e., blue-green algae). Examples of suitable eukaryotic host cells include, for example, mammalian cells, plant cells, yeast cells such as *Saccharomyces cerevisiae*, *Saccharomyces carlsbergensis*, *Lipomyces starkey*, *Candida* spp. such as *Yarrowia (Candida) lipolytica*, *Kluyveromyces* spp., *Pichia* spp., *Trichoderma* spp. or *Hansenula* spp., or fungal cells such as filamentous fungal cells, for

example, *Aspergillus*, *Neurospora* and *Penicillium*. Preferably, *Saccharomyces cerevisiae* (baker's yeast) cells are utilized.

Expression in a host cell can be accomplished in a transient or stable fashion. Transient expression can occur from introduced constructs which contain expression signals functional in the host cell, but which constructs do not replicate and rarely integrate into the host cell, or where the host cell is not proliferating. Transient expression also can be accomplished by inducing the activity of a regulatable promoter operably linked to the gene of interest, although such inducible systems frequently exhibit a low basal level of expression. Stable expression can be achieved by introducing a construct that can integrate into the host genome or that autonomously replicates in the host cell. Stable expression of the gene of interest can be selected through the use of a selectable marker located on or co-transfected with the expression construct, followed by selection for cells expressing the marker. When stable expression results from integration, the site of the construct's integration can occur randomly within the host genome or can be targeted through the use of constructs containing regions of homology with the host genome sufficient to target recombination with the host locus. Where constructs are targeted to an endogenous locus, all or some of the transcriptional and translational regulatory regions can be provided by the endogenous locus.

A transgenic mammal may also be used in order to express the enzymes of the present invention, and thus ultimately produce the PUFA(s) of interest. More specifically, once the above-described construct is created, it may be inserted into the pronucleus of an embryo. The embryo may then be implanted into a recipient female. Alternatively, a nuclear transfer method can also be utilized (Schnieke et al., Science 278:2130-2133 (1997)). Gestation and birth are then permitted to occur (see, e.g., U.S. Patent No. 5,750,176 and U.S. Patent No. 5,700,671). Milk, tissue, or other fluid samples from the offspring should then contain altered levels of PUFAs, as compared to the levels

normally found in the non-transgenic animal. Subsequent generations may be monitored for production of the altered or enhanced levels of PUFAs and thus incorporation of the gene(s) encoding the desired desaturase enzyme(s) into their genomes. The mammal utilized as the host may be selected from the group consisting of, for example, mice, rats, rabbits, swine (porcines), goats and sheep (ovines), horses, and bovines. However, any mammal may be used provided it has the ability to incorporate DNA encoding the enzyme of interest into its genome.

For expression of a desaturase polypeptide, functional transcriptional and translational initiation and termination regions are operably linked to the DNA encoding the desaturase polypeptide of interest. Transcriptional and translational initiation and termination regions are derived from a variety of nonexclusive sources, including the DNA to be expressed, genes known or suspected to be capable of expression in the desired system, expression vectors, chemical synthesis, or from an endogenous locus in a host cell. Expression in a plant tissue and/or plant part presents certain efficiencies, particularly where the tissue or part is one which is harvested early, such as seed, leaves, fruits, flowers, roots, etc. Expression can be targeted to that location with the plant by utilizing specific regulatory sequence such as those of U.S. Patent Nos. 5,463,174; 4,943,674; 5,106,739; 5,175,095; 5,420,034; 5,188,958; and 5,589,379.

Alternatively, the expressed protein can be an enzyme that produces a product, and that product may be incorporated, either directly or upon further modifications, into a fluid fraction from the host plant. Expression of a desaturase gene, or antisense desaturase transcripts, can alter the levels of specific PUFAs, or derivatives thereof, found in plant parts and/or plant tissues. The desaturase polypeptide coding region may be expressed either by itself or with other genes, in order to produce tissues and/or plant parts containing higher proportions of desired PUFAs, or in which the PUFA composition

more closely resembles that of human breast milk (Prieto et al., PCT publication WO 95/24494).

The termination region may be derived from the 3' region of the gene from which the initiation region was obtained or from a different gene. A large number of termination regions are known to be satisfactory in a variety of hosts from the same and different genera and species. The termination region usually is selected as a matter of convenience rather than because of any particular property.

As noted above, a plant (e.g., *Glycine max* (soybean) or *Brassica napus* (canola)) or plant tissue may also be utilized as a host or host cell, respectively, for expression of the desaturase enzymes, which may, in turn, be utilized in the production of PUFAs. More specifically, desired PUFAs can be expressed in seed. Methods of isolating seed oils are known in the art. Thus, in addition to providing a source for PUFAs, seed oil components may be manipulated through the expression of the desaturase genes, as well as perhaps other desaturase genes and elongase genes, to provide seed oils that can be added to nutritional compositions, pharmaceutical compositions, animal feeds and cosmetics. Once again, a vector that comprises a DNA sequence encoding the desaturase gene of interest, operably linked to a promoter, is introduced into the plant tissue or plant for a time and under conditions sufficient for expression of the desaturase gene. The vector may also comprise one or more genes that encode other enzymes, for example, delta-5-desaturase, elongase, delta-12-desaturase, delta-15-desaturase, delta-17-desaturase, and/or delta-19-desaturase enzymes. The plant tissue or plant may produce the relevant substrate (e.g., adrenic acid or DPA) upon which the enzyme acts or a vector encoding enzymes that produce such substrates may be introduced into the plant tissue, plant cell or plant. In addition, suitable substrates may be sprayed on plant tissues expressing the appropriate enzymes. Using these various techniques, one may produce PUFAs by use of a plant cell, plant tissue, or plant. It should also

be noted that the invention also encompasses a transgenic plant comprising the above-described vector, wherein expression of the nucleotide sequence(s) of the vector results in production of a desired PUFA in, for example, the seeds of the transgenic plant.

The regeneration, development, and cultivation of plants from single plant protoplast transformants or from various transformed explants is well known in the art (Weissbach and Weissbach, In: Methods for Plant Molecular Biology, (Eds.), Academic Press, Inc. San Diego, CA, (1988)). This regeneration and growth process typically includes the steps of selection of transformed cells, culturing those individualized cells through the usual stages of embryonic development through the rooted plantlet stage. Transgenic embryos and seeds are similarly regenerated. The resulting transgenic rooted shoots are thereafter planted in an appropriate plant growth medium such as soil.

The development or regeneration of plants containing the foreign, exogenous gene that encodes a protein of interest is well known in the art. Preferably, the regenerated plants are self-pollinated to provide homozygous transgenic plants. Otherwise, pollen obtained from the regenerated plants is crossed to seed-grown plants of agronomically important lines. Conversely, pollen from plants of these important lines is used to pollinate regenerated plants. A transgenic plant of the present invention containing a desired polypeptide is cultivated using methods well known to one skilled in the art.

There are a variety of methods for the regeneration of plants from plant tissue. The particular method of regeneration will depend on the starting plant tissue and the particular plant species to be regenerated.

Methods for transforming dicots, primarily by use of *Agrobacterium tumefaciens*, and obtaining transgenic plants have been published for cotton (U.S. Patent No. 5,004,863, U.S. Patent No. 5,159,135, U.S. Patent No. 5,518, 908); soybean (U.S. Patent No. 5,569,834, U.S. Patent No. 5,416,011, McCabe et. al., *BiolTechnology* 6:923 (1988), Christou et al., *Plant Physiol.*

87:671-674 (1988)); Brassica (U.S. Patent No. 5,463,174); peanut (Cheng et al., Plant Cell Rep. 15:653-657 (1996), McKently et al., Plant Cell Rep. 14:699-703 (1995)); papaya; and pea (Grant et al., Plant Cell Rep. 15:254-258, (1995)).

Transformation of monocotyledons using electroporation, particle bombardment, and Agrobacterium have also been reported. Transformation and plant regeneration have been achieved in asparagus (Bytebier et al., Proc. Natl. Acad. Sci. (USA) 84:5354, (1987)); barley (Wan and Lemaux, Plant Physiol 104:37 (1994)); Zea mays (Rhodes et al., Science 240:204 (1988), Gordon-Kamm et al., Plant Cell 2:603-618 (1990), Fromm et al., BiolTechnology 8:833 (1990), Kozziel et al., BiolTechnology 11: 194, (1993), Armstrong et al., Crop Science 35:550-557 (1995)); oat (Somers et al., BiolTechnology 10: 15 89 (1992)); orchard grass (Horn et al., Plant Cell Rep. 7:469 (1988)); rice (Toriyama et al., TheorAppl. Genet. 205:34, (1986); Part et al., Plant Mol. Biol. 32:1135-1148, (1996); Abedinia et al., Aust. J. Plant Physiol. 24:133-141 (1997); Zhang and Wu, Theor. Appl. Genet. 76:835 (1988); Zhang et al. Plant Cell Rep. 7:379, (1988); Batraw and Hall, Plant Sci. 86:191-202 (1992); Christou et al., Bio/Technology 9:957 (1991)); rye (De la Pena et al., Nature 325:274 (1987)); sugarcane (Bower and Birch, Plant J. 2:409 (1992)); tall fescue (Wang et al., BiolTechnology 10:691 (1992)), and wheat (Vasil et al., Bio/Technology 10:667 (1992); U.S. Patent No. 5,631,152).

Assays for gene expression based on the transient expression of cloned nucleic acid constructs have been developed by introducing the nucleic acid molecules into plant cells by polyethylene glycol treatment, electroporation, or particle bombardment (Marcotte et al., Nature 335:454-457 (1988); Marcotte et al., Plant Cell 1:523-532 (1989); McCarty et al., Cell 66:895-905 (1991); Hattori et al., Genes Dev. 6:609-618 (1992); Goff et al., EMBO J. 9:2517-2522 (1990)).

Transient expression systems may be used to functionally dissect gene constructs (see generally, Maliga et al., Methods in Plant Molecular Biology, Cold Spring Harbor Press (1995)). It is

understood that any of the nucleic acid molecules of the present invention can be introduced into a plant cell in a permanent or transient manner in combination with other genetic elements such as vectors, promoters, enhancers etc.

In addition to the above discussed procedures, practitioners are familiar with the standard resource materials which describe specific conditions and procedures for the construction, manipulation and isolation of macromolecules (e.g., DNA molecules, plasmids, etc.), generation of recombinant organisms and the screening and isolating of clones, (see for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press (1989); Maliga et al., *Methods in Plant Molecular Biology*, Cold Spring Harbor Press (1995); Birren et al., *Genome Analysis: Detecting Genes*, 1, Cold Spring Harbor, New York (1998); Birren et al., *Genome Analysis: Analyzing DNA*, 2, Cold Spring Harbor, New York (1998); *Plant Molecular Biology: A Laboratory Manual*, eds. Clark, Springer, New York (1997)).

In view of the above, the present invention encompasses a method of producing an omega-3 desaturase and/or a delta-12 desaturase enzyme, the method comprising the steps of: 1) isolating the nucleotide sequence of the gene encoding the desired desaturase enzyme(s); 2) constructing a vector comprising the nucleotide sequence(s); and 3) introducing the vector into a host cell for a time and under conditions sufficient for the production of the desaturase enzyme(s).

The present invention also encompasses a method of producing PUFAs, the method comprising exposing a suitable fatty acid substrate to the enzyme such that the desaturase converts the fatty acid substrate to a desired PUFA product. For example, when AA (20:4n-6) is exposed to the omega-3 desaturase enzyme of the present invention, it is converted into EPA (20:5n-3). The EPA so formed may be converted into DPA (22:5n-3) by the action of an elongase, and the DPA subsequently converted into DHA (22:6n-3) by a delta-4 desaturase.

Likewise, when OA (18:1n-9) is exposed to the delta-12 desaturase enzyme of the present invention, it is converted into

LA (18:2n-6). The LA so formed may be converted into virtually all of the PUFAs shown in Figure 1 by the subsequent actions of suitable desaturases and/or elongases.

Uses of the Subject Desaturase Genes and Enzymes Encoded Thereby:

As noted above, the isolated desaturase genes and the desaturase enzymes encoded thereby have many uses. For example, the genes and the corresponding enzymes may be used indirectly or directly, singly or in combination, in the production of PUFAs. For example, the omega-3 desaturase may be used in the production of ETA, EPA, DPA, DHA, and the like. As used in this context, the word "directly" encompasses the situation where the enzyme is used to catalyze the conversion of a fatty acid substrate directly into the desired fatty acid product, without any intermediate steps or pathway intermediates (e.g., the conversion of AA to EPA). The product so obtained is then utilized in a composition. "Indirectly" encompasses the situation where a desaturase according to the present invention is used to catalyze the conversion of a fatty acid substrate into another fatty acid (i.e., a pathway intermediate) by the desaturase (e.g., the conversion of AA to EPA) and then the latter fatty acid (the EPA) is converted to the desired fatty acid product by use of another desaturase or non-desaturase enzyme (e.g., the conversion of EPA to DPA by elongase). These PUFAs (i.e., those produced either directly or indirectly by the activity of the subject desaturases) may be added to, for example, nutritional compositions, pharmaceutical compositions, cosmetics, and animal feeds, all of which are encompassed by the present invention. Such uses are described in detail below.

Nutritional Compositions:

The present invention includes nutritional compositions. For purposes of the present invention, such compositions include any food or preparation for human consumption (including for enteral and/or parenteral consumption) which when taken into the body (a)

serve to nourish or build up tissues or supply energy; and/or (b) maintain, restore or support adequate nutritional status or metabolic function.

The nutritional composition of the present invention comprises an oil, fatty acid ester, or fatty acid produced directly or indirectly by use of the desaturase genes disclosed herein. The composition may either be in a solid or liquid form. Additionally, the composition may include edible macronutrients, vitamins, and/or minerals in amounts desired for a particular use. The amounts of these ingredients will vary depending on whether the composition is intended for use with normal, healthy infants, children, or adults, or for use with individuals having specialized needs, such as individuals suffering from metabolic disorders and the like.

Examples of macronutrients that may be added to the compositions include (but are not limited to): edible fats, carbohydrates and proteins. Examples of such edible fats include (but are not limited to): coconut oil, borage oil, fungal oil, black current oil, soy oil, and mono- and diglycerides. Examples of such carbohydrates include (but are not limited to): glucose, edible lactose, and hydrolyzed starch. Additionally, examples of proteins which may be utilized in the nutritional composition of the invention include (but are not limited to) soy proteins, electrodialysed whey, electrodialysed skim milk, milk whey, or the hydrolysates of these proteins.

With respect to vitamins and minerals, the following may be added to the nutritional compositions of the present invention: calcium, phosphorus, potassium, sodium, chloride, magnesium, manganese, iron, copper, zinc, selenium, iodine, and Vitamins A, E, D, C, and the B complex. Other such vitamins and minerals may also be added.

The components utilized in the nutritional compositions of the present invention will be of semi-purified or purified origin. By semi-purified or purified is meant a material which

has been prepared by purification of a natural material or by de novo synthesis.

Examples of nutritional compositions of the present invention include (but are not limited to): infant formulas, dietary supplements, dietary substitutes, and rehydration compositions. Nutritional compositions of particular interest include (but are not limited to) compositions for enteral and parenteral supplementation for infants, specialized infant formulas, supplements for the elderly, and supplements for those with gastrointestinal difficulties and/or malabsorption.

The nutritional composition of the present invention may also be added to food even when supplementation of the diet is not required. For example, the composition may be added to food of any type, including (but not limited to): margarine, modified butter, cheeses, milk, yogurt, chocolate, candy, snacks, salad oils, cooking oils, cooking fats, meats, fish and beverages.

In a preferred embodiment of the present invention, the nutritional composition is an enteral nutritional product, more preferably, an adult or pediatric enteral nutritional product. This composition may be administered to adults or children experiencing stress or having specialized needs due to chronic or acute disease states. The composition may comprise, in addition to PUFAs produced according to the present invention, macronutrients, vitamins, and/or minerals, as described previously. The macronutrients may be present in amounts equivalent to those present in human milk or on an energy basis, i.e., on a per calorie basis.

The enteral formula, for example, may be sterilized and subsequently utilized on a ready-to-feed (RTF) basis or stored in a concentrated liquid or powder. The powder can be prepared by spray-drying the formula prepared as indicated above, and reconstituting it by rehydrating the concentrate. Adult and pediatric nutritional formulas are known in the art and are commercially available (e.g., Similac®, Ensure®, Jevity® and Alimentum® from Ross Products Division, Abbott Laboratories,

Columbus, Ohio). An oil or fatty acid produced in accordance with the present invention may be added to any of these formulas.

The energy density of the nutritional compositions of the present invention, when in liquid form, may range from about 0.6 kcal to about 3 kcal per ml. When in solid or powdered form, the nutritional supplements may contain from about 1.2 to more than 9 kcals per gram, preferably about 3 to 7 kcals per gm. In general, the osmolality of a liquid product should be less than 700 mOsm and, more preferably, less than 660 mOsm.

The nutritional formula may include macronutrients, vitamins, and minerals, as noted above, in addition to the PUFAs produced in accordance with the present invention. The presence of these additional components helps the individual ingest the minimum daily requirements of these elements. In addition to the provision of PUFAs, it may also be desirable to add zinc, copper, folic acid, and antioxidants to the composition. It is believed that these substances boost a stressed immune system and will therefore provide further benefits to the individual receiving the composition. A pharmaceutical composition may also be supplemented with these elements.

In a more preferred embodiment, the nutritional composition comprises, in addition to antioxidants and at least one PUFA, a source of carbohydrate wherein at least 5 weight percent of the carbohydrate is indigestible oligosaccharide. In a more preferred embodiment, the nutritional composition additionally comprises protein, taurine, and carnitine.

As noted above, the PUFAs produced in accordance with the present invention, or derivatives thereof, may be added to a dietary substitute or supplement, particularly an infant formula, for patients undergoing intravenous feeding or for preventing or treating malnutrition or other conditions or disease states. As background, it should be noted that human breast milk has a fatty acid profile comprising from about 0.15% to about 0.36% DHA, from about 0.03% to about 0.13% EPA, from about 0.30% to about 0.88% AA, from about 0.22% to about 0.67% DGLA, and from about 0.27% to about 1.04% GLA. Thus, fatty acids such as AA, EPA and/or DHA

produced in accordance with the present invention can be used to alter, for example, the composition of infant formulas in order to replicate more faithfully the PUFA content of human breast milk or to alter the presence of PUFAs normally found in a non-human mammal's milk. In particular, a composition for use as a medicinal agent or food supplement, particularly a breast milk substitute or supplement, will preferably comprise one or more of AA, DGLA and GLA. More preferably, the composition will comprise from about 0.3 to 30% AA, from about 0.2 to 30% DGLA, and/or from about 0.2 to about 30% GLA.

Parenteral nutritional compositions comprising from about 2 to about 30% by weight fatty acids calculated as triglycerides are encompassed by the present invention. The preferred composition has about 1 to about 25% by weight of the total PUFA composition as GLA. Other vitamins, particularly fat-soluble vitamins such as vitamin A, D, E and L-carnitine, can optionally be included. When desired, a preservative such as alpha-tocopherol may be added in an amount of about 0.1% by weight.

In addition, the ratios of AA, DGLA and GLA can be adapted for a particular given end use. When formulated as a breast milk supplement or substitute, a composition which comprises one or more of AA, DGLA and GLA will be provided in a ratio of from about 1:19:30 to about 6:1:0.2, respectively. For example, the breast milk of animals can vary in ratios of AA:DGLA:GLA ranging from 1:19:30 to 6:1:0.2, which includes intermediate ratios which are preferably about 1:1:1, 1:2:1, 1:1:4. When produced together in a host cell, adjusting the rate and percent of conversion of a precursor substrate such as GLA and DGLA to AA can be used to control the PUFA ratios precisely. For example, a 5% to 10% conversion rate of DGLA to AA can be used to produce an AA to DGLA ratio of about 1:19, whereas a conversion rate of about 75% to 80% can be used to produce an AA to DGLA ratio of about 6:1. Therefore, whether in a cell culture system or in a host animal, regulating the timing, extent and specificity of desaturase expression, as well as the expression of other desaturases and elongases, can be used to modulate PUFA levels and ratios. The

PUFAs produced in accordance with the present invention (e.g., AA, EPA, etc.) may then be combined with other PUFAs or other types of fatty acids in the desired concentrations and ratios.

Additionally, PUFAs produced in accordance with the present invention or host cells transformed to contain and express the subject desaturase genes may also be used as animal food supplements to alter an animal's tissue or milk fatty acid composition to one more desirable for human or animal consumption.

Pharmaceutical Compositions:

The present invention also encompasses a pharmaceutical composition comprising one or more of the acids and/or resulting oils produced using the desaturase genes, in accordance with the methods described herein. Specifically, such a pharmaceutical composition may comprise one or more of the PUFAs and/or oils, in combination with a standard, well-known, non-toxic pharmaceutically-acceptable carrier, adjuvant or vehicle such as phosphate-buffered saline, water, ethanol, polyols, vegetable oils, a wetting agent or an emulsion such as a water/oil emulsion. The composition may be in either a liquid or solid form. For example, the composition may be in the form of a tablet, capsule, ingestible liquid or powder, injectible, or topical ointment or cream. Proper fluidity can be maintained, for example, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants. It may also be desirable to include isotonic agents, for example, sugars, sodium chloride, and the like. Besides such inert diluents, the composition can also include adjuvants, such as wetting agents, emulsifying and suspending agents, sweetening agents, flavoring agents and perfuming agents.

Suspensions, in addition to the active compounds, may comprise suspending agents such as ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar, and tragacanth or mixtures of these substances.

Solid dosage forms such as tablets and capsules can be prepared using techniques well known in the art. For example, PUFAs produced in accordance with the present invention can be tableted with conventional tablet bases such as lactose, sucrose, and cornstarch in combination with binders such as acacia, cornstarch or gelatin, disintegrating agents such as potato starch or alginic acid, and a lubricant such as stearic acid or magnesium stearate. Capsules can be prepared by incorporating these excipients into a gelatin capsule along with antioxidants and the relevant PUFA(s). The antioxidant and PUFA components should fit within the guidelines presented above.

For intravenous administration, the PUFAs produced in accordance with the present invention or derivatives thereof may be incorporated into commercial formulations such as Intralipids™. The typical normal adult plasma fatty acid profile comprises 6.64 to 9.46% AA, 1.45 to 3.11% DGLA, and 0.02 to 0.08% GLA. These PUFAs or their metabolic precursors can be administered alone or in combination with other PUFAs to achieve a normal fatty acid profile in a patient. Where desired, the individual components of the formulations may be provided individually, or in kit form, for single or multiple use. A typical dosage of a particular fatty acid is from 0.1 mg to 20 g, taken from one to five times per day (up to 100 g daily) and is preferably in the range of from about 10 mg to about 1, 2, 5, or 10 g daily (taken in one or multiple doses).

Possible routes of administration of the pharmaceutical compositions of the present invention include, for example, enteral (e.g., oral and rectal) and parenteral. For example, a liquid preparation may be administered orally or rectally. Additionally, a homogenous mixture can be completely dispersed in water, admixed under sterile conditions with physiologically acceptable diluents, preservatives, buffers or propellants to form a spray or inhalant.

The route of administration will, of course, depend upon the desired effect. For example, if the composition is being utilized to treat rough, dry, or aging skin, to treat injured or

burned skin, or to treat skin or hair affected by a disease or condition, it may be applied topically.

The dosage of the composition to be administered to the patient may be determined by one of ordinary skill in the art and depends upon various factors such as weight of the patient, age of the patient, overall health of the patient, past history of the patient, immune status of the patient, etc.

With respect to form, the composition may be, for example, a solution, a dispersion, a suspension, an emulsion or a sterile powder that is then reconstituted.

The present invention also includes the treatment of various disorders by use of the pharmaceutical and/or nutritional compositions described herein. In particular, the compositions of the present invention may be used to treat restenosis after angioplasty. Furthermore, symptoms of inflammation, rheumatoid arthritis, asthma and psoriasis may also be treated with the compositions of the invention. Evidence also indicates that PUFAs may be involved in calcium metabolism; thus, the compositions of the present invention may be utilized in the treatment or prevention of osteoporosis and of kidney or urinary tract stones.

Additionally, the compositions of the present invention may also be used in the treatment of cancer. Malignant cells have been shown to have altered fatty acid compositions. Addition of fatty acids has been shown to slow their growth, cause cell death and increase their susceptibility to chemotherapeutic agents. Moreover, the compositions of the present invention may also be useful for treating cachexia associated with cancer.

The compositions of the present invention may also be used to treat diabetes (see U.S. Patent No. 4,826,877 and Horrobin et al., Am. J. Clin. Nutr. Vol. 57 (Suppl.) 732S-737S). Altered fatty acid metabolism and composition have been demonstrated in diabetic animals.

Furthermore, the compositions of the present invention, comprising PUFAs produced either directly or indirectly through the use of the desaturase enzymes, may also be used in the

treatment of eczema and in the reduction of blood pressure. Additionally, the compositions of the present invention may be used to inhibit platelet aggregation, to induce vasodilation, to reduce cholesterol levels, to inhibit proliferation of vessel wall smooth muscle and fibrous tissue (Brenner et al., Adv. Exp. Med. Biol. Vol. 83, p.85-101, 1976), to reduce or to prevent gastrointestinal bleeding and other side effects of non-steroidal anti-inflammatory drugs (see U.S. Patent No. 4,666,701), to prevent or to treat endometriosis and premenstrual syndrome (see U.S. Patent No. 4,758,592), and to treat myalgic encephalomyelitis and chronic fatigue after viral infections (see U.S. Patent No. 5,116,871).

Further uses of the compositions of the present invention include use in the treatment of AIDS, multiple sclerosis, and inflammatory skin disorders, as well as for maintenance of general health.

Additionally, the composition of the present invention may be utilized for cosmetic purposes. It may be added to pre-existing cosmetic compositions such that a mixture is formed or a PUFA produced according to the subject invention may be used as the sole "active" ingredient in a cosmetic composition.

Veterinary Applications:

It should be noted that the above-described pharmaceutical and nutritional compositions may be utilized in connection with animals (i.e., domestic or non-domestic, including mammals, birds, reptiles, lizards, etc.), as well as humans, as animals experience many of the same needs and conditions as humans. For example, the oil or fatty acids of the present invention may be utilized in animal feed supplements, animal feed substitutes, animal vitamins or in animal topical ointments.

The present invention may be further illustrated by the non-limiting examples presented below:

EXAMPLE 1

Construction of *Saprolegnia diclina* (ATCC 56851) cDNA Library

To isolate genes encoding for functional desaturase enzymes, a cDNA library was constructed. *Saprolegnia diclina* cultures were grown in potato dextrose media (Difco # 336, BD Diagnostic Systems, Sparks, Maryland) at room temperature for four days with constant agitation. The mycelia were harvested by filtration through several layers of cheesecloth, and the cultures were crushed in liquid nitrogen using a mortar and pestle. The cell lysates were resuspended in RT buffer (Qiagen, Valencia, California) containing β -mercaptoethanol and incubated at 55°C for three minutes. These lysates were homogenized either by repeated aspirations through a syringe or over a "Qiashredder"-brand column (Qiagen). The total RNA was finally purified using the "RNeasy Maxi"-brand kit (Qiagen), as per the manufacturer's protocol.

mRNA was isolated from total RNA from each organism using an oligo dT cellulose resin. The "pBluescript II XR"-brand library construction kit (Stratagene, La Jolla, California) was used to synthesize double-stranded cDNA. The double-stranded cDNA was then directionally cloned (5' *EcoRI*/3' *XhoI*) into pBluescript II SK(+) vector (Stratagene). The *S. diclina* library contained approximately 2.5×10^6 clones, each with an average insert size of approximately 700 bp. Genomic DNA of *S. diclina* was isolated by crushing the culture in liquid nitrogen followed by purification using the "Genomic DNA Extraction"-brand kit (Qiagen), as per the manufacturer's protocol.

EXAMPLE 2

Determination of Codon Usage in *Saprolegnia diclina*

The 5' ends of 350 random cDNA clones were sequenced from the *Saprolegnia diclina* cDNA library described in Example 1. The sequences were translated into six reading frames using GCG

program (Genetics Computer Group, Madison, Wisconsin) with the ``FastA''-brand algorithm to search for similarity between a query sequence and a group of sequences of the same type, specifically within the GenBank database. Many of the clones were identified as putative housekeeping genes based on protein homology to known genes. Eight *S. diclina* cDNA sequences were thus selected. Additionally, the full-length *S. diclina* delta 5-desaturase and delta 6-desaturase sequences were also used (see Table 1 below). These sequences were then used to generate the *S. diclina* codon bias table shown in Table 2 below by employing the ``CodonFrequency'' program from GCG.

Table 1
Genes from *Saprolegnia diclina* used for
generation of Codon Bias Table

Clone #	Match	# bases	# amino acids
3	Actin gene	615	205
20	Ribosomal protein L23	420	140
55	Heat Shock protein 70 gene	468	156
83	Glyceraldehyde-3-P-dehydrogenase gene	588	196
138	Ribosomal protein S13 gene	329	110
179	Alpha-tubulin 3 gene	591	197
190	Casein kinase II alpha subunit gene	627	209
250	Cyclophilin gene	489	163
	Delta 6-desaturase	1362	453
	Delta 5-desaturase	1413	471
	Total	6573	2191

Table 2
Codon Bias Table for *Saprolegnia diclina*

Amino acid	Codon Bias	% used
Ala	GCC	55%
Arg	CGC	50%
Asn	AAC	94%
Asp	GAC	85%
Cys	TGC	77%
Gln	CAG	90%
Glu	GAG	80%
Gly	GGC	67%
His	CAC	86%
Ile	ATC	82%
Leu	CTC	52%
Lys	AAG	87%
Met	ATG	100%
Phe	TTC	72%
Pro	CCG	55%
Ser	TCG	47%
Thr	ACG	46%
Trp	TGG	100%
Tyr	TAC	90%
Val	GTC	73%
Stop	TGA	67%

EXAMPLE 3

Design of Degenerate Oligonucleotides for the Isolation of an
Omega-3 Desaturase from *Saprolegnia diclina* (ATCC 56851)

Fungi like *Saprolegnia diclina* produce a wide range of PUFAs, including arachidonic acid (AA) and eicosapentaenoic acid

(EPA) via the PUFA biosynthetic pathway depicted in Figure 1. Analysis of the fatty acid composition of *Saprolegnia diclina* (ATCC 56851) showed 15.42% of the total lipid to be AA and 12.2% of the total lipid to be EPA (see Table 5). Linoleic acid (LA) was the only other intermediate present in high amounts. This indicates that *S. diclina* has very active delta-6 and delta-5 desaturases, as well as elongases that shunt intermediates through the pathway depicted in Figure 1. Due to the high percentage of EPA in this organism, an active omega-3 desaturase (synonymous with a ``delta-15'' desaturase when the substrate is a C₁₈ fatty acid, a ``delta-17'' desaturase when the substrate is a C₂₀ fatty acid, and a ``delta-19'' desaturase when the substrate is a C₂₂ fatty acid) is predicted to exist which is capable of converting AA (20:4n-6) to EPA (20:5n-3).

As just noted, omega-3 desaturases are enzymes that catalyze the introduction of a double bond at the delta-15 position for C₁₈-acyl chains, the delta-17 position for C₂₀-acyl chains, and the delta-19 position for C₂₂-acyl chains. There are several known omega-3 desaturases from plants, but these act exclusively on C₁₈ fatty acid substrates like LA (18:2n-6) and GLA (18:3n-6). These types of desaturases are collectively referred to as delta 15-desaturases. At this point in time, only one omega-3 desaturase gene has been isolated whose encoded enzyme catalyzes the desaturation of C₁₈, C₂₀, and C₂₂ fatty acid substrates. This is *fat-1* from *C. elegans*. See U.S. Patent No. 6,194,167, issued February 27, 2001.

The approach used in this Example to identify an omega-3 desaturase from *S. diclina* involved PCR amplification of a region of the desaturase gene using degenerate oligonucleotides (primers) that contained conserved motifs present in other known omega-3 desaturases. Omega-3 desaturases from the following organisms were used for the design of these degenerate primers: *Arabidopsis thaliana* (Swissprot # P46310), *Ricinus communis* (Swissprot # P48619), *Glycine max* (Swissprot # P48621), *Sesamum*

indicum (Swissprot # P48620), *Nicotiana tabacum* (GenBank # D79979), *Perilla frutescens* (GenBank # U59477), *Capsicum annuum* (GenBank # AF222989), *Limnanthes douglassi* (GenBank # U17063), and *Caenorhabditis elegans* (GenBank # L41807). Some primers were designed to contain the conserved histidine-box motifs that are important components of the active site of the enzymes. See Shanklin, J.E., McDonough, V.M., and Martin, C.E. (1994) *Biochemistry* 33, 12787-12794.

Alignment of sequences was carried out using the CLUSTALW Multiple Sequence Alignment Program (<http://workbench.sdsc.edu>).

The following degenerate primers were designed and used in various combinations:

Protein Motif 1: NH₃- TRAAIPKHCWVK -COOH

Primer RO 1144 (Forward): 5'-ATC CGC GCC GCC ATC CCC AAG CAC TGC TGG GTC AAG-3' (SEQ ID NO: 1).

Protein Motif 2: NH₃- ALFVLGHDCGHGSFS -COOH

This primer contains the histidine-box 1 (underlined).

Primer RO 1119 (Forward): 5'- GCC CTC TTC GTC CTC GGC CAY GAC TGC GGC CAY GGC TCG TTC TCG-3' (SEQ. ID. NO: 2).

Primer RO 1118 (Reverse): 5'-GAG RTG GTA RTG GGG GAT CTG GGG GAA GAR RTG RTG GRY GAC RTG-3' (SEQ. ID. NO: 3).

Protein Motif 3: NH₃- PYHGWRISHRTHHQON -COOH

This primer contains the histidine-box 2 (underlined).

Primer RO 1121 (Forward): 5'-CCC TAC CAY GGC TGG CGC ATC TCG CAY CGC ACC CAY CAY CAG AAC-3' (SEQ. ID. NO: 4).

Primer RO 1122 (Reverse): 5'-GTT CTG RTG RTG GGT CCG RTG CGA GAT GCG CCA GCC RTG GTA GGG-3' (SEQ. ID. NO: 5).

Protein Motif 4: NH₃- GSHF D/H P D/Y SDLFV -COOH

Primer RO 1146 (Forward): 5'- GGC TCG CAC TTC SAC CCC KAC
TCG GAC CTC TTC GTC-3' (SEQ. ID. NO: 6).

Primer RO 1147 (Reverse): 5'- GAC GAA GAG GTC CGA GTM GGG
GTW GAA GTG CGA GCC-3' (SEQ. ID. NO: 7).

Protein Motif 5: NH₃- WS Y/F L/V RGGLTT L/I DR -COOH

Primer RO 1148 (Reverse): 5'- GCG CTG GAK GGT GGT GAG GCC
GCC GCG GAW GSA CGA CCA-3' (SEQ. ID. NO: 8).

Protein Motif 6: NH₃- HHDIGTHVIIHHLLFPQ -COOH

This sequence contains the third histidine-box (underlined).

Primer RO 1114 (Reverse): 5'- CTG GGG GAA GAG RTG RTG GAT
GAC RTG GGT GCC GAT GTC RTG RTG-3' (SEQ. ID. NO: 9).

Protein Motif 7: NH₃- H L/F FP Q/K IPHYHL V/I EAT -COOH

Primer RO 1116 (Reverse): 5'- GGT GGC CTC GAY GAG RTG GTA
RTG GGG GAT CTK GGG GAA GAR RTG-3' (SEQ. ID. NO: 10).

Protein Motif 8: NH₃- HV A/I HH L/F FPQIPHYHL -COOH

This primer contains the third histidine-box (underlined)
and accounts for differences between the plant omega-3
desaturases and the *C. elegans* omega-3-desaturase.

Primer RO 1118 (Reverse): 5'-GAG RTG GTA RTG GGG GAT CTG GGG
GAA GAR RTG RTG GRY GAC RTG-3' (SEQ. ID. NO: 11).

The degeneracy code used for SEQ. ID. NOS: 1 through 11 was
as follows: R= A/G; Y=C/T; M=A/C; K=G/T; W=A/T; S=C/G; B=C/G/T;
D=A/G/T; H=A/C/T; V=A/C/G; and N=A/C/G/T.

EXAMPLE 4Identification and Isolation of an Omega-3 Desaturase Gene from *Saprolegnia diclina* (ATCC 56851)

Various sets of the degenerate primers disclosed in Example 3 were used in PCR amplification reactions, using as a template either the *S. diclina* cDNA library plasmid DNA (from Example 1), or *S. diclina* genomic DNA. Also various different DNA polymerases and reaction conditions were used for the PCR amplifications. These reactions variously involved using "Platinum Taq"-brand DNA polymerase (Life Technologies Inc., Rockville, Maryland), or cDNA polymerase (Clontech, Palo Alto, California), or Taq PCR-mix (Qiagen), at differing annealing temperatures.

PCR amplification using the primers RO 1121 (Forward) (SEQ. ID. NO: 4) and RO 1116 (Reverse) (SEQ. ID. NO: 10) resulted in the successful amplification of a fragment homologous to a known omega-3 desaturase. The RO 1121 (Forward) primer corresponds to the protein motif 3; the RO 1116 (Reverse) primer corresponds to the protein motif 7.

PCR amplification was carried out in a 50 µl total volume containing: 3 µl of the cDNA library template, PCR buffer containing 40 mM Tricine-KOH (pH 9.2), 15 mM KOAc, 3.5 mM Mg(OAc)₂, 3.75 µg/ml BSA (final concentration), 200 µM each deoxyribonucleotide triphosphate, 10 pmole of each primer and 0.5 µl of "Advantage"-brand cDNA polymerase (Clontech). Amplification was carried out as follows: initial denaturation at 94°C for 3 minutes, followed by 35 cycles of the following: 94°C for 1 min, 60°C for 30 sec, 72°C for 1 min. A final extension cycle of 72°C for 7 min was carried out, followed by reaction termination at 4°C.

A single ~480 bp PCR band was generated which was resolved on a 1% "SeaKem Gold"-brand agarose gel (FMC BioProducts, Rockland, Maine), and gel-purified using the Qiagen Gel

Extraction Kit. The staggered ends on the fragment were ``filled-in'' using T4 DNA polymerase (Life Technologies, Rockville, Maryland) as per the manufacturer's instructions, and the DNA fragments were cloned into the PCR-Blunt vector (Invitrogen, Carlsbad, California). The recombinant plasmids were transformed into TOP10 supercompetent cells (Invitrogen), and eight clones were sequenced and a database search (Gen-EMBL) was carried out.

Clones ``sdd17-7-1'' to ``sdd17-7-8' were all found to contain and ~483 bp insert. The deduced amino acid sequence from this fragment showed highest identity to the following proteins (based on a ``tFastA'' search):

1. 37.9% identity in 161 amino acid overlap with an omega-3 (delta-15) desaturase from *Synechocystis* sp. (Accession # D13780).

2. 40.7% identity in 113 amino acid overlap with *Picea abies* plastidial omega-3 desaturase (Accession # AJ302017).

3. 35.9% identity in 128 amino acid overlap with *Zea mays* FAD8 fatty acid desaturase (Accession # D63953).

Based on its sequence homology to known omega-3 fatty acid desaturases, it seemed likely that this DNA fragment was part of an omega-3 desaturase gene present in *S. diclina*.

The DNA sequence identified above was used in the design oligonucleotides to isolate the 3' and the 5' ends of this gene from the cDNA library described in Example 1. To isolate the 3' end of the gene, the following oligonucleotides were designed:

RO 1188 (Forward): 5'-TAC GCG TAC CTC ACG TAC TCG CTC G-3' (SEQ. ID. NO: 12).

RO 1189 (Forward): 5'-TTC TTG CAC CAC AAC GAC GAA GCG ACG-3' (SEQ. ID. NO: 13).

RO 1190 (Forward): 5'-GGA GTG GAC GTA CGT CAA GGG CAA C-3' (SEQ. ID. NO: 14).

RO 1191 (Forward): 5'-TCA AGG GCA ACC TCT CGA GCG TCG AC-3' (SEQ. ID. NO: 15).

These primers (SEQ. ID. NOS: 12-15) were used in combinations with the pBluescript SK(+) vector oligonucleotide: RO 898: 5'-CCC AGT CAC GAC GTT GTA AAA CGA CGG CCA G-3' (SEQ. ID. NO: 16).

PCR amplifications were carried out using either the "Taq PCR Master Mix" brand polymerase (Qiagen) or "Advantage"-brand cDNA polymerase (Clontech) or "Platinum"-brand Taq DNA polymerase (Life Technologies), as follows:

For the "Taq PCR Master Mix" polymerase, 10 pmoles of each primer were used along with 1 μ l of the cDNA library DNA from Example 1. Amplification was carried out as follows: initial denaturation at 94°C for 3 min, followed by 35 cycles of the following: 94°C for 1 min, 60°C for 30 sec, 72°C for 1 min. A final extension cycle of 72°C for 7 min was carried out, followed by the reaction termination at 4°C. This amplification resulted in the most distinct bands as compared with the other two conditions tested.

For the "Advantage"-brand cDNA polymerase reaction, PCR amplification was carried out in a 50 μ l total volume containing: 1 μ l of the cDNA library template from Example 1, PCR buffer containing 40 mM Tricine-KOH (pH 9.2), 15 mM KOAc, 3.5 mM Mg(OAc)₂, 3.75 μ g/ml BSA (final concentration), 200 μ M each deoxyribonucleotide triphosphate, 10 pmole of each primer and 0.5 μ l of cDNA polymerase (Clontech). Amplification was carried out as described for the Taq PCR Master Mix.

For the "Platinum"-brand Taq DNA polymerase reaction, PCR amplification was carried out in a 50 μ l total volume containing: 1 μ l of the cDNA library template from Example 1, PCR buffer containing 20 mM Tris-Cl, pH 8.4, 50 mM KCl (final concentration), 200 μ M each deoxyribonucleotide triphosphate, 10 pmole of each primer, 1.5 mM MgSO₄, and 0.5 μ l of Platinum Taq DNA polymerase. Amplification was carried out as follows:

initial denaturation at 94°C for 3 min, followed by 30 cycles of the following: 94°C for 45 sec, 55°C for 30 sec, 68°C for 2 min. The reaction was terminated at 4°C.

All four sets of primers in combination with the vector primer generated distinct bands. PCR bands from the combination (RO 1188 + RO 898) were >500 bp and this was gel-purified and cloned separately. The PCR bands generated from the other primer combinations were <500 bp. The bands were gel-purified, pooled together, and cloned into PCR-Blunt vector (Invitrogen) as described earlier. The recombinant plasmids were transformed into TOP10 supercompetent cells (Invitrogen) and clones were sequenced and analyzed.

Clone ``sdd17-16-4'' and ``sdd16-6'' containing the ~500 bp insert, and clones ``sdd17-17-6,'' ``sdd17-17-10,'' and ``sdd17-20-3,'' containing the ~ 400 bp inserts, were all found to contain the 3'-end of the putative omega-3 desaturase. These sequences overlapped with each other, as well as with the originally identified fragment of this putative omega-3 desaturase gene. All of the sequences contained the 'TAA' stop codon and a poly-A tail typical of 3'-ends of eukaryotic genes. This 3'-end sequence was homologous to other known omega-3 desaturases, sharing the highest identity (41.5% in 130 amino acid overlap) with the *Synechocystis* delta-15 desaturase (Accession # D13780).

For the isolation of the 5'-end of the this gene, the following oligonucleotides were designed and used in combinations with the pBluescript SK(+) vector oligonucleotide:

RO 899: 5'- AGC GGA TAA CAA TTT CAC ACA GGA AAC AGC -3'
(SEQ. ID. NO: 17).

RO 1185 (Reverse): 5'-GGT AAA AGA TCT CGT CCT TGT CGA TGT TGC-3' (SEQ. ID. NO: 18).

RO 1186 (Reverse): 5'-GTC AAA GTG GCT CAT CGT GC-3' (SEQ. ID. NO: 19).

RO 1187 (Reverse): 5'-CGA GCG AGT ACG TGA GGT ACG CGT AC-3'
(SEQ. ID. NO: 20).

Amplifications were carried out using either the ``Taq PCR Master Mix''-brand polymerase (Qiagen) or the ``Advantage''-brand cDNA polymerase (Clontech) or the ``Platinum''-brand Taq DNA polymerase (Life Technologies), as described hereinabove for the 3' end isolation.

PCR bands generated from primer combinations (RO 1185 or RO 1186 + RO 899) were between ~580 to ~440 bp and these were pooled and cloned into PCR-Blunt vector as described above. Clones thus generated included ``sdd17-14-1,'' ``sdd17-14-10,'' ``sdd17-18-2,'' and ``sdd17-18-8,'' all of which showed homology with known omega-3 desaturases.

Additionally, bands generated from (RO 1187 + RO 899) were ~680 bp, and these were cloned separately to generate clones ``sdd17-22-2'' and ``sdd17-22-5'' which also showed homology with known omega-3 desaturases. All these clones overlapped with each other, as well as with the original fragment of the unknown putative omega-3 desaturase. These sequences contained an ATG' site followed by an open reading frame, indicating that it is the start site of this gene. These fragments showed highest identity (39.7% in 146 amino acid overlap) with the delta-15 desaturase from *Calendula officinalis* (Accession # AJ245938).

The full-length of this omega-3 desaturase was obtained by PCR amplification of the *S. diclina* cDNA library using the following oligonucleotides:

RO 1212 (Forward): 5'-TCA ACA GAA TTC ATG ACC GAG GAT AAG ACG AAG GTC GAG TTC CCG-3' (SEQ. ID. NO: 21). This primer contains the 'ATG' start site (single underline) followed by the 5' sequence of the omega-3 desaturase. In addition, an *EcoRI* site (double underline) was introduced upstream of the start site to facilitate cloning into the yeast expression vector pYX242.

RO 1213 (Reverse): 5'-AAA AGA AAG CTT CGC TTC CTA GTC TTA GTC CGA CTT GGC CTT GGC-3' (SEQ. ID. NO: 22). This primer contains the 'TAA' stop codon (single underline) of the gene as well as sequence downstream from the stop codon. This sequence was included because regions within the gene were very G+C rich,

and thus could not be included in the design of oligonucleotides for PCR amplification. In addition, a *HindIII* site (double underline) was included for convenient cloning into a yeast expression vector pYX242.

PCR amplification was carried out using the "Taq PCR Master Mix"-brand polymerase (Qiagen), 10 pmoles of each primer, and 1 µl of the cDNA library DNA from Example 1. Amplification was carried out as follows: initial denaturation at 94°C for 3 min, followed by 35 cycles of the following: 94°C for 1 min, 60°C for 30 sec, 72°C for 1 min. A final extension cycle of 72°C for 7 min was carried out, followed by the reaction termination at 4°C.

A PCR band of ~1 kb was thus obtained and this band was isolated, purified, cloned into PCR-Blunt vector (Invitrogen), and transformed into TOP10 cells. The inserts were sequenced to verify the gene sequence. Clone "sdd17-27-2" was digested with *EcoRI* / *HindIII* to release the full-length insert, and this insert was cloned into yeast expression vector pYX242, previously digested with *EcoRI* / *HindIII*. This construct contained 1077 bp of sdd17 cloned into pYX242. This construct was labeled pRSP19, which was transformed into yeast SC334 for expression studies.

In addition, the *S. diclina* omega-3 gene was cloned into another yeast expression vector, pYES2 (Invitrogen). For this, the omega-3 desaturase gene was isolated from the cDNA library generated in Example 1 by PCR amplification (as described above) using the following oligonucleotides:

RO1221 (Forward) (SEQ. ID. NO: 23)

5'-TCA ACA AAG CTT ATG ACC GAG GAT AAG ACG AAG GTC GAG TTC CCG-3'

This primer contains the 'ATG' start site (underlined) followed by the 5' sequence of the omega-3 desaturase. In addition, a *HindIII* site (bold) was introduced upstream of the start site to facilitate cloning into the pYES2 yeast expression vector.

RO1222 (Reverse) (SEQ. ID. NO: 24)

5'-AAA AGA GAA TTC CGC TTC CTA GTC TTA GTC CGA CTT GGC CTT GGC-3'

This primer contains the 'TAA' stop codon (underlined) of the gene as well as sequence downstream from the stop codon. This sequence was included since regions within the gene were very G+C rich, and thus could not be included in the design of oligonucleotides for PCR amplification. In addition, an *EcoRI* site (bold) was included for convenient cloning into the pYES2 yeast expression vector.

The ~1 kb PCR band thus generated was digested with *HindIII*/*EcoRI*, and cloned into pYES2 digested with the same restriction enzymes. The resulting construct (*sdd17* + pYES2) was labeled pRSP20, and was used in co-expression studies.

Attempts were also made to isolate the full-length *sdd17* gene from genomic DNA by PCR amplification. However, the PCR product obtained was larger than 1077 bp (~ 1.15 kb), and sequencing of this product revealed the presence of small introns in the genomic sequence.

The full-length gene of this putative omega-3 desaturase (labeled *sdd17*) was 1077 bp in length and is shown in Figure 2 (SEQ ID NO: 25).

The gene of SEQ ID NO: 25 encoded a protein of 358 amino acid residues (SEQ. ID. NO: 26) (Figure 3). A search of the deduced protein sequence of *sdd17* (using the 'tFastA' program) showed the protein to have highest identity (41% in 269 amino acid overlap) with the delta-15 desaturase from *Synechocystis* sp. (ATCC Accession No. 13780) (Figure 4) and *Synechocystis* sp. PCC6803 (ATCC Accession No. D90913). This protein shared sequence similarities with several other plant omega-3 desaturases. Comparison of this predicted protein sequence with the FAT1 enzyme from *C. elegans* (ATCC Accession L41807) revealed only a 31.6% identity in 310 amino acid overlap (Figure 5).

Like all omega-3 desaturases, this enzyme does not contain a cytochrome b5 domain within the 5' end of its sequence. The cytochrome b5 domain is present in most 'front-end' desaturating enzymes like delta 5- and delta 6-desaturases. The

omega-3 desaturase described in this example includes the three histidine-rich sequences that are present in all membrane-bound desaturases. These three domains are present at position 89 to 93 (HDCGH), 125 to 129 (HRHHH), and 284 to 288 (HQVHH) of SEQ. ID. NO: 26. These histidine-rich boxes are believed to coordinate the diiron-oxo structure at the enzyme's active site, and are necessary for enzyme activity; see Stuke, J.E., McDonough, V.M. & Martn, C.E. (1990) *J. Biol. Chem.* 265, 20144-20149. These features are consistent with the "SDD17" protein being a member of the membrane-bound desaturase /hydroxylase family of the diiron-oxo proteins. The G+C content of this gene is 61.8%.

EXAMPLE 5

Expression of the Omega-3 Desaturase Gene ("sdd17") from *Saprolegnia diclina* in Bakers' Yeast

To determine the substrate specificity and the class of reaction catalyzed by the SDD17-protein, *sdd17* was heterologously expressed in a *Saccharomyces cerevisiae* (SC334). Because *S. cerevisiae* cannot synthesize fatty acids beyond OA (18:1n-9), it is an ideal system to use to determine enzyme activity on substrates longer than OA because no background enzyme activity will be detected. Suitable fatty acid substrates can be exogenously supplied to the host which are taken up by the cell and acted on by the expressed protein of the transformed *sdd17* gene.

Clone pRSP19, which contained the full-length omega-3 desaturase (*sdd17*) from *S. diclina* cloned into pYX242, was transformed into *Saccharomyces cerevisiae* (SC334) using the "Alkali-Cation Yeast Transformation"-brand kit (BIO 101, Vista, California). Transformants were selected for leucine auxotrophy on media lacking leucine (DOB [-Leu]). To detect the specific desaturase activity of these clones, transformants were grown in the presence of 50 μ M of each of LA (18:2n-6), GLA (18:3n-6),

DGLA (20:3n-6), AA (20:4n-6), and adrenic acid (22:4n-6). Conversion of these exogenously supplied fatty acid substrates into a product having one additional unsaturation indicates the presence of a specific desaturase activity that is not found in the wild-type *S. cerevisiae*:

Conversion of LA (18:2n-6) to ALA (18:3n-3) indicates delta-15 desaturase activity.

Conversion of GLA (18:3n-6) to STA (18:4n-3) indicates delta-15 desaturase activity.

Conversion of DGLA (20:3n-6) to ETA (20:4n-3) indicates delta-17 desaturase activity.

Conversion of AA (20:4n-6) to EPA (20:5n-3) indicates delta-17 desaturase activity.

Conversion of adrenic acid (22:4n-6) to DPA (22:5n-3) indicates delta-19 desaturase activity.

The negative control strain was *S. cerevisiae* transformed with the pYX242 vector. The experimental cultures and the control cultures were grown simultaneously and analyzed.

The cultures were vigorously agitated (250 rpm) and grown for 48 hours at 24°C in the presence of 50 µM (final concentration) of the various substrates (see Table 3). The cells were spun down, washed once in distilled water, and the pellets resuspended in methanol; chloroform was added along with tritridecanoin (as an internal standard). These mixtures were incubated for at least an hour at room temperature, or at 4°C overnight. The chloroform layer was extracted and filtered through a Whatman filter with 1 gm anhydrous sodium sulfate to remove particulate matter and residual water. The organic solvents were evaporated at 40°C under a stream of nitrogen. The extracted lipids were then converted to fatty acid methyl esters (FAME) for gas chromatography analysis (GC) by adding 2 ml 0.5 N potassium hydroxide in methanol to a closed tube. The samples were heated to 95°C-100°C for 30 minutes and cooled to room

temperature. Approximately 2 ml 14% borontrifluoride in methanol was added and the heating repeated. After the extracted lipid mixture cooled, 2 ml of water and 1 ml of hexane were added to extract the FAME for analysis by GC. The percent conversion was calculated using the formula:

$$\% \text{ Conversion} = \frac{[\% \text{ Product}]}{[\% \text{ Product} + \% \text{ Substrate}]} \times 100$$

Table 3
Yeast Expression of the Omega-3 Desaturase (SDD17) from *Saprolegnia diclina* at 24°C

Clone	Substrate* Incorporated	Product Produced	% Conversion	Enzyme Activity
PRSP19	18:2 n-6 (9.42%)	18:3 n-3 (0%)	0	Delta 15
	18:3 n-6 (9.11%)	18:4 n-3 (0%)	0	Delta 15
	20:3 n-6 (21.36%)	20:4 n-3 (1.18%)	5%	Delta 17
	20:4 n-6 (32.14%)	20:5 n-3 (5.16%)	13.8%	Delta 17
	22:4 n-6 (28.65%)	22:5 n-3 (1.22%)	4%	Delta 19
Control (pYX242)	18:2 n-6 (9.27%)	18:3 n-3 (0%)	0	Delta 15
	18:3 n-6 (9.18%)	18:4 n-3 (0%)	0	Delta 15
	20:3 n-6 (14.19%)	20:4 n-3 (0%)	0	Delta 17
	20:4 n-6 (26.56%)	20:5 n-3 (0%)	0	Delta 17
	22:4 n-6 (16.4%)	22:5 n-3 (0%)	0	Delta 19

* 50 μ M substrate used

Numbers in parenthesis represent fatty acid as a percentage of total lipids from yeast.

18:2n-6 = Linoleic acid	20:3n-6= dihomogamma-linolenic acid
18:3n-6= gamma-linolenic acid	20:4n-6= arachidonic acid
18:3n-3= alpha-linolenic acid	22:4n-6= adrenic acid
18:4n-3= stearidonic acid	20:4n-3= eicosatetraenoic acid
20:5n-3= eicosapentaenoic acid	22:5n-3= omega-3-docosapentaenoic acid

Table 3 displays the enzyme activity of the *sdd17*-encoded protein product from *Saprolegnia diclina* (ATCC 56851). This enzyme is an active omega-3 desaturase capable of desaturating both C₂₀ and C₂₂ omega-6 fatty acid substrates to yield the corresponding omega-3 fatty acid products. This enzyme converted 13.8% of the added AA substrate to the corresponding EPA product, thus indicating delta-17 desaturase activity. In addition, this enzyme also acted on DGLA, converting it to ETA, as would be expected for a delta-17 desaturase. In this Example, however, only 5% of the added DGLA was converted to ETA, indicating that under the conditions used here, the enzyme has a substrate preference for AA as compared to DGLA.

The activity of this enzyme toward C₂₂ fatty acid substrates was also investigated because C₂₂ omega-3 fatty acids like DPA and DHA have important dietary and pharmaceutical implications. From Table 3, it can be seen that this enzyme was active on C₂₂ substrates such as adrenic acid, converting 4% of it to DPA. As can be seen from the control cultures, there was no non-specific conversion of exogenously added substrate to product in non-transformed *S. cerevisiae*.

Table 4 demonstrates that this omega-3 desaturase (SDD17) can also function at a lower temperature. (i.e. 15°C). Here, 50

μM of exogenous substrate was added to the transformants and the cultures were grown for 48 hours at 15°C . Fatty acid analysis was carried out as described above. The overall uptake of substrate by *S. cerevisiae* at 15°C was lower than that seen at 24°C (compare Table 3 & Table 4). However the percent conversion of substrate to product by the enzyme increased at 15°C . Since the presence of lower concentration of exogenous fatty acid substrate seemed to improve the activity of the enzyme, it is possible that fatty acid substrates at high concentrations may exert a feed-back inhibition on this enzyme. Further studies may be carried out to determine the effect of different substrate concentration and different temperatures on expression of this omega-3 desaturase in *S. cerevisiae*.

Table 4
Yeast Expression of the Omega-3 Desaturase (SDD17) from *Saprolegnia*
diclina at 15°C

Clone	Substrate* Incorporated	Product Produced	% Conversion	Enzyme Activity
PRSP19	18:2 n-6 (8.79%)	18:3 n-3 (0%)	0	Delta 15
	18:3 n-6 (12.69%)	18:4 n-3 (0%)	0	Delta 15
	20:3 n-6 (9.52%)	20:4 n-3 (1.46%)	13%	Delta 17
	20:4 n-6 (8.69%)	20:5 n-3 (2.05%)	19%	Delta 17
	22:4 n-6 (6.68%)	22:5 n-3 (0.62%)	8.4%	Delta 19
Control (pYX242)	18:2 n-6 (9.65%)	18:3 n-3 (0%)	0	Delta 15
	18:3 n-6 (13.55%)	18:4 n-3 (0%)	0	Delta 15
	20:3 n-6 (10.17%)	20:4 n-3 (0%)	0	Delta 17
	20:4 n-6 (16.58%)	20:5 n-3 (0%)	0	Delta 17
	22:4 n-6 (11.05%)	22:5 n-3 (0%)	0	Delta 19

* 50 μ M substrate used

Numbers in parenthesis represent fatty acid as a percentage of total lipids from yeast

18:2n-6 = Linoleic acid

18:3n-6= gamma-linolenic acid

18:3n-3= alpha-linolenic acid

18:4n-3= stearidonic acid

20:5n-3= eicosapentaenoic acid

20:3n-6= dihomogamma-linolenic acid

20:4n-6= arachidonic acid

22:4n-6= adrenic acid

20:4n-3= eicosatetraenoic acid

22:5n-3= omega-3-docosapentaenoic acid

Unlike all known omega-3 desaturases, the *sdd17*-encoding enzyme did not desaturate any C₁₈ omega-6 fatty acyl substrates to their corresponding omega-3 fatty acids (under the conditions tested). It is possible that *in vivo*, this enzyme functions exclusively on AA, converting it to EPA. This would be consistent with the fatty acid profile of *S. diclina* displaying high amounts of AA and EPA, but little or none of the other omega-3 intermediates (Table 5).

Table 5

Fatty Acid Profile of *Saprolegnia diclina* ATCC 56851

Fatty Acid	% Total Lipid
C10:0	0.22
C12:0	0.1
C13:0	3.98
C14:0	6.0
C14:1 n-5	0.29
C15:0	1.06
C16:0	19.75
C16:1 n-7	1.99
C18:0	3.99
C18: n-9	15.39
C18:1 n-7	7.39
C18:1 n-5	0.43
C18:2 n-6	7.07
C18:3 n-6	2.13
C18:3 n-3	0.08
C20:0	0.76
C20:1 n-9	0.15
C20:1 n-7	0.08
C20:2 n-6	0.22
C20:3 n-6	1.31
C20:4 n-6	15.42
C20:5n-3	12.2

Thus, *sdd17* encodes a novel omega-3 desaturase, capable of desaturating C₂₀ and C₂₂ fatty acid substrates. The SDD17 protein can easily be expressed in a heterologous system and thus has potential for use in other heterologous systems like plants. This enzyme is very different from other known omega-3

desaturases, showing activity on both C₂₀ and C₂₂ fatty acid substrates, but not C₁₈ substrates. It shares only 31.6% identity with FAT-1, the only other known desaturase capable of acting on C₂₀ and C₂₂ omega-6 fatty acid substrates. Thus the enzyme encoded by *sdd17* is a novel omega-3 desaturase.

EXAMPLE 6

Co-Expression of *S. diclina* Omega-3 Desaturase with Other Enzymes

The pRSP20 construct consisting of *sdd17* cloned into pYES2 yeast expression vector, as described in Example 3, was used in co-expression studies with other desaturases and elongases. pRSP3, a construct that contained the delta 5-desaturase gene (SEQ ID NO: 27) from *S. diclina* cloned into the pYX242 yeast expression vector, was co-transformed with pRSP20 into yeast. Transformation protocol was as described in Example 4. This delta 5-desaturase catalyzes the conversion of DGLA to AA and ETA to EPA. Co-transformants were selected on minimal media lacking leucine and Uracil (DOB [-Leu-Ura]).

Table 6 shows that when 50 μ M of the substrate DGLA (20:3 n-6) was added, the delta 5-desaturase converted it to AA (20:4, n-6), and the omega-3 desaturase was able to further desaturate AA to EPA (24:5, n-3). The percent conversion of the substrate to final product was 5%, with no background observed in the negative control.

Table 6
Co-expression Studies with the Omega-3 Desaturase (SDD17)
from *S. diclina*

Clone	Plasmid in yeast	20:3 n-6 (DGLA) Incorporated	20:4 n-6 (AA) Produced	20:5 n-3 (EPA) Produced	% Conversion
Cntrl	pYX242 + pYES2	19.33	0	0	0
pRSP22	pRSP3 (Delta 5) + pRSP20 (omega-3 desaturase)	20.56	2.64	0.14	5%
Clone	Plasmid in yeast	18:3 n-6 (GLA) Incorporated	20:3 n-6 (DGLA) Produced	20:4 n-3 (ETA) Produced	% Conversion
Cntrl	pYX242 + pYES2	4.83	0	0	0
pRSP23	pRAT-4-A7 (elongase) + pRSP20 (omega-3 desaturase)	4.56	9.30	0.14	1.4%

* 50 μ M substrate used

Numbers represent fatty acid as a percentage of total lipids from yeast

18:3n-6= gamma-linolenic acid

20:3n-6= dihomo-gamma-linolenic acid

20:4n-6= arachidonic acid

20:4n-3= eicosatetraenoic acid

20:5n-3= eicosapentaenoic acid

$$\% \text{ Conversion} = \frac{[\% \text{ Product 1} + \% \text{ Product 2}]}{[\% \text{ substrate} + \% \text{ Product 1} + \% \text{ Product 2}]}$$

Table 6 also shows the results of a co-transformation experiment involving pRSP20 and pRAT-4-A7, an elongase obtained from *Thraustochytrid* sp. 7091 (SEQ. ID. NO: 28) cloned into pYX242. This elongase gene catalyzes the addition of two more carbons to the pre-existing PUFA. When 50 μ M of the substrate GLA (18:3 n-6) was added to the co-transformants, the elongase elongated GLA to DGLA, and the DGLA was further desaturated by the omega-3 desaturase to ETA (20:4 n-3). The percent conversion of substrate to final product was 1.4%, with no background observed in the negative control.

Thus the *S. diclina* omega-3 desaturase was able to utilize a product produced, in a heterologous expression system, by another heterologous enzyme from the PUFA biosynthetic pathway, and convert that product to the expected PUFA.

It should be noted that the expression (and hence the activity) of *sdd17* when cloned in the pYES2 vector (pRSP20) was much lower than when cloned into the pYX242 vector (pRSP19). This could be explained by the difference in the expression promoters present in each vector. The pYX242 promoter is a constitutive promoter and is much stronger than the galactose-inducible promoter in pYES2. Similar observations have been made during expression studies with other desaturases when cloned into these two expression vectors.

Further co-expression studies may be carried out using pRSP19 instead of pRSP20 along with other desaturases and elongases. Also the *S. diclina* omega-3 desaturase may also be co-expressed with other enzymes like the delta 4-desaturase pRTA7 (SEQ. ID. NO: 29), where in adrenic acid (22:4 n-6) may be added as a substrate and the final end product DHA (22:6 n-3) may be produced due to the concerted action of the omega-3 desaturase and the delta 4-desaturase.

EXAMPLE 7Design of Degenerate Primers for the Isolation of the Delta 12-Desaturase Gene from *Saprolegnia diclina* ATCC 56851

Analysis of the fatty acid composition of *Saprolegnia diclina* (ATCC 56851) revealed the presence of a considerable amount of LA, which suggested the presence of a very active delta 12-desaturase (Table 5). Delta 12-desaturases use OA as a substrate, thus catalyzing the conversion of OA to LA (see Figure 1). Delta 12-desaturases are present only in plants, fungi, and insects, but not in mammals, including humans. Thus LA is an "essential" fatty acid in humans because it cannot be synthesized *in vivo*. LA is further desaturated and elongated to produce important intracellular compounds like GLA, AA, and EPA.

The goal of this experiment was to isolate the delta 12-desaturase gene from *S. diclina* and verify its functionality by expressing the enzyme in a heterologous host system such as yeast. The approach taken was to design degenerate primers (oligonucleotides) that represent conserved amino acid motifs from known delta 12-desaturases. In designing these primers, known delta-12 desaturase sequence information from both fungi and plant sources was used, including sequence information from: *Mortierella alpina* (Accession #AF110509), *Mucor rouxii* (Accession #AF161219), *Brassica juncea* (Accession #X91139), *Arabidopsis thaliana* (Accession #L26296), and *Borago officinalis* (Accession #AF0744324). The sequence information was analyzed using the CODEHOP Blockmaker program (<http://blocks.fhcrc.org/codehop.html>).

The degenerate primers used in this Example were as follows:

Protein Motif 1: NH₃- P N/E FTIKEIR D/E A/C IPAHC F -COOH

Primer RO 967 (Forward): 5'-CCG SAG TTC ACS ATC AAG GAG ATC CGC GAS KSC ATC CCG GCC CAC TGC TTC -3' (SEQ. ID. NO: 30).

Protein Motif 2: NH₃- MP H/F YHAEET V/Y H I/L KK A/L -COOH

Primer RO 968 (Reverse): 5'-GRS CTT CTT GAK GTG GWM SGT GGC CTC CTC GGC GTG GTA GWR CGG CAT-3' (SEQ. ID. NO: 31).

Protein Motif 3: NH₂- P L/V YW A/I C/M/A QG V/I V L/G/C TGVW -COOH

Primer RO 964 (Forward): 5'-CCS STC TAC TGG GCC TGC CAG GGT RTC GTC CTC ACS GGT GTC TGG-3' (SEQ. ID. NO: 32).

This sequence is more similar to the known plant Delta 12-desaturases.

Primer RO 965 (Forward): 5'-CCS STC TAC TGG ATC RYS CAG GGT RTC GTC KGY ACS GGT GTC TGG-3' (SEQ. ID. NO: 33).

This sequence is more similar to the known fungal Delta 12-desaturases.

Protein Motif 4: NH₂- HVAHH L/F FS T/Q MPHYHA -COOH

Primer RO 966 (Reverse): 5'-GGC GTG GTA GTG CGG CAT SMM CGA GAA GAR GTG GTG GGC GAC GTG-3' (SEQ. ID. NO: 34).

The degeneracy code used for the oligonucleotides was as follows: R= A/G; Y=C/T; M=A/C; K=G/T; W=A/T; S=C/G; B=C/G/T; D=A/G/T; H=A/C/T; V=A/C/G; N=A/C/G/T.

EXAMPLE 8

Identification and Isolation of the Delta 12-Desaturase Gene from *Saprolegnia diclina* (ATCC 56851)

To isolate a fragment of the delta 12-desaturase gene from *S. diclina*, PCR was carried out using the *S. diclina* cDNA library from Example 1 as a template. Primers were used in the following combinations: (RO 964 + RO 966), (RO 965 + RO 966), and (RO 967 + RO 968). PCR was carried out in 100 µl volumes using the "Taq PCR Master Mix"-brand polymerase (Qiagen). 10 pmoles of each primer were used along with 1 µl of the cDNA library DNA. Amplification was carried out as follows: initial denaturation at 94°C for 4 min, followed by 25 cycles of the following: 94°C for 1 min, 47°C for 1 min, 72°C for 2 min. A final extension cycle of

72°C for 5 min was carried out, followed by reaction termination at 4°C.

Amplification with (RO 964 + RO 966) or (RO 965 + RO 966) resulted in distinct bands of ~688 bp in length. Amplification with (RO 967 + RO 968) resulted in one distinct band of ~660 bp. These bands were resolved on a 1% "SeaKem Gold"-brand agarose gel (FMC BioProducts), and gel-purified using the Qiagen Gel Extraction Kit. The staggered ends on the fragment were "filled-in" using T4 DNA polymerase (Life Technologies), following the manufacturer's specifications. The DNA fragments were then cloned into the PCR-Blunt vector (Invitrogen). The recombinant plasmids were transformed into TOP10 supercompetent cells (Invitrogen), clones were sequenced, and a database search (Gen-EMBL) was carried out.

Clones "sdd12-1-8," "sdd12-2-8," and "sdd12-5-1" were all found to overlap with each other, and these overlapping fragments were aligned using the "ASSEMBLE"-brand program (GCG) to create a single fusion fragment of ~900 bp. A "tFastA" search with the deduced amino acids of this fusion sequence showed highest identity to the following proteins:

49% identity in 298 amino acid overlap with *Borago officinalis* Delta 12-desaturase (Accession #AF074324) and 46.7% identity in 332 amino acid overlap with *Sesamum indicum* Delta 12-desaturase (Accession # AF192486).

Based on the high identity to known delta 12-desaturases, the fragment was considered to be a region of the *S. diclina* delta 12-desaturase gene. This fragment was used to design primers to pull up the 5'- and 3'-ends of the gene.

To isolate the 3' end of the gene, the following oligonucleotides were designed:

RO 975 (Forward): 5'-CAC GTA CCT CCA GCA CAC GGA CAC CTA CG-3' (SEQ. ID. NO: 35).

RO 976 (Forward): 5'- GAT CGA CAG CGC GAT CCA CCA CAT TGC-3' (SEQ. ID. NO: 36).

These were used in combinations with the pBluescript SK(+) vector oligonucleotide RO 898: 5'-CCC AGT CAC GAC GTT GTA AAA CGA CGG CCA G-3' (SEQ. ID. NO: 16).

PCR amplifications were carried out using ``Taq PCR Master Mix''-brand polymerase (Qiagen) as follows: 10 pmoles of each primer were used along with 1 µl of the cDNA library DNA from Example 1 as template. Amplification conditions were as follows: initial denaturation at 94°C for 3 min, followed by 35 cycles of the following: 94°C for 1 min, 60°C for 30 sec, 72°C for 1 min. A final extension cycle of 72°C for 7 min was carried out, followed by reaction termination at 4°C.

Primer combination (RO 898 + RO 975) generated a PCR band of ~390 bp and primer combination (RO 898 + RO 976) generated a band of length ~300 bp. These bands were purified and cloned into PCR-Blunt vector as described earlier. Several clones, including clones ``sdd12-8-12'' and ``sdd12-9-4'' were found to contain the 3' end of the delta 12-desaturase gene. These sequences overlapped the initial delta-12 desaturase fragment and included a TAA stop codon and a poly-A tail. Sequence analysis with the ``tFastA'' program revealed that clone ``sdd12-9-4'' shared 54.5% identity in a 73 amino acid overlap with the delta 12-desaturase from *Mortierella alpina* (Accession #AB020033), and 56.9% identity in a 72 amino acid overlap with the delta 12-desaturase from *Mucor rouxii* (Accession #AF161219).

To isolate the 5' end of the this gene, the following oligonucleotides were designed and used in combinations with the pBluescript SK(+) vector oligonucleotide RO 899 (SEQ. ID. NO: 17).

RO 977 (Reverse): 5'- CAA ATG GTA AAA GCT AGT GGC AGC GCT GC-3' (SEQ ID NO: 37).

RO 978 (Reverse): 5'-AGT ACG TGC CCT GGA CGA ACC AGT AGA TG-3' (SEQ ID NO: 38).

PCR amplifications were carried out using either ``Taq PCR Master Mix''-brand polymerase (Qiagen) or the ``Advantage-GC''-

brand cDNA PCR kit (Clonetech). The Clonetech product was used to circumvent potential PCR amplification problems that may occur with GC-rich regions generally present at the 5'-end of desaturases from this organism. PCR amplifications using the ``Taq PCR Master Mix''-brand polymerase was carried out as described for the isolation of 3'-end of this gene.

When using the ``Advantage-GC cDNA PCR''-brand kit, thermocycling conditions were as follows: the template was initially denatured at 94°C for 1 min, followed by 30 cycles of 94°C for 30 seconds, 68°C for 3 minutes, and finally an extension cycle at 68°C for 5 min. Each reaction included 1 µl of cDNA library template (from Example 1), 10 pmole of each primer, 0.2 mM each dNTP, 1M GC Melt, 40 mM Tricine-KOH, 15 mM KOAc, 3.5 mM MG(OAc)₂, 5% DMSO, and 375 µg/ml BSA in a total volume of 50 µl.

A PCR product of ~371 bp was obtained using the primer combination (RO 899 + RO 978). This band was cloned into the PCR-Blunt vector (Invitrogen) as described earlier. Only one clone, ``sdd12-10-8,'' thus obtained contained the putative ATG start site of the gene. Other clones had the ATG replaced by other codons. ``tFastA'' analysis of the deduced amino acid sequence of ``sdd12-10-8'' showed 47.2% identity in a 72 amino acid overlap of the delta-12 desaturase from *Impatiens balsamina* (Accession #AF182520) and 42.7% identity in a 75 amino acid overlap with the delta-12 desaturase from *Calendula officinalis* (Accession #AJ245938).

The full length of this delta 12-desaturase was obtained by PCR amplification of the *S. diclina* cDNA library of Example 1 using the following oligonucleotides:

RO1051 (Forward): 5'- TCA ACA GAA TTC ATG TGC AAA GGT CAA GCT CCT TCC AAG GCC GAC GTG -3' (SEQ. ID. 39). This primer contains the `ATG' start site (underlined) followed by the 5' sequence of the Delta 12-desaturase. In addition, an *EcoRI* site (double underline) was introduced upstream of the start site to facilitate cloning into the pYX242 yeast expression vector.

RO1057 (Reverse): 5'- AAA AGA AAG CTT TTA CTT TTC CTC GAG CTT GCG CTT GTA AAA CAC AAC-3' (SEQ. ID. NO: 40). This primer contains the TAA stop codon (underlined) of the gene as well as a *HindIII* site (double underline), which was included for convenient cloning into the pYX242 yeast expression vector.

PCR amplifications were carried out using both "Taq PCR Master Mix"-brand polymerase (Qiagen) and the "Advantage-GC cDNA PCR"-brand kit (Clonetech), as described earlier. In this case, however, *S. diclina* genomic DNA was used as the template for amplification. A PCR band of ~1.1 kb was thus obtained and this band was isolated, purified, cloned into PCR-Blunt vector (Invitrogen), and transformed into TOP10 cells. The inserts were sequenced to verify the gene sequence. Clone "sdd12-gg-b1" was digested with *EcoRI* / *HindIII* to release the full-length insert, and this insert was cloned into the yeast expression vector pYX242, previously digested with *EcoRI* / *HindIII*. This construct included 1182 bp of the delta-12 desaturase gene and pYX242. The construct was labeled pRSP11. The pRSP11 construct was then transformed into *S. cerevisiae* (SC334) for expression studies.

The full-length gene of this putative delta-12 desaturase (labeled *sdd12*) was 1182 bp in length (SEQ ID NO: 41) (Figure 6). The gene encodes a protein of 393 amino acid residues (SEQ ID NO: 42) (Figure 7). A "tFastA" search of the deduced protein sequence of *sdd12* showed the protein to have highest identity (45.6% in a 379 amino acid overlap) with the delta-12 desaturase from *Gossypium hirsutum* (Accession #X97016) (Figure 8) and 49.6% identity in a 353 amino acid overlap with the delta-12 desaturase from *Sesamum indicum* (FAD2) (Accession #AF192486).

Like other delta-12 desaturases, this enzyme also does not contain a cytochrome b5 domain within the 5' end of its' sequence. This enzyme does contain the three histidine-rich sequences that are present in all membrane-bound desaturases. The position and length of these histidine-boxes are similar to those seen in other desaturases. These are present at amino acid

positions 108 to 112 (HECGH), 144 to 148 (HRRHH), and 326 to 330 (HVTHH) of SEQ. ID. NO: 42. As noted earlier, these histidine-rich boxes are believed to co-ordinate the diiron-oxo structure at the enzyme's active site and are necessary for enzyme activity.

EXAMPLE 9

Expression of the Delta 12-Desaturase Gene (*sdd12*) in Bakers' Yeast

To determine the substrate specificity and the class of reaction catalyzed by this delta 12-desaturase (SDD12), *sdd12* was heterologously expressed in a *Saccharomyces cerevisiae* (SC334). As noted earlier, because *S. cerevisiae* cannot synthesize fatty acids beyond OA, it is an ideal system to determine enzyme activity on substrates longer than OA because no background enzyme activity will be detected. Suitable fatty acid substrates are exogenously supplied to the host; these substrates are taken up by the cell and acted on by the expressed delta-12 desaturase of the transformed *sdd12* gene.

Clone PRSP11, which contained the full-length delta-12 desaturase (*sdd12*) from *S. diclina*, cloned into pYX242, was transformed into *Saccharomyces cerevisiae* (SC334) using the "Alkali-Cation Yeast Transformation"-brand kit (BIO 101), following the manufacturer's instructions. Transformants were selected for leucine auxotrophy on media lacking leucine (DOB-Leu). To detect the specific desaturase activity of these clones, transformants were grown in the presence of 50 μ M each of OA, LA, GLA, and DGLA.

Conversion of OA to LA (18:2n-6) indicates delta-12 desaturase activity.

Conversion of LA to ALA (18:3n-3) indicates delta-15 desaturase activity.

Conversion of LA to GLA (18:3n-6) indicates delta-6 desaturase activity.

Conversion of GLA to stearidonic (18:4n-3) acid indicates delta-15 desaturase activity.

Conversion of DGLA to ETA (20:4n-3) indicates delta-17 desaturase activity.

Conversion of DGLA to AA (20:4n-6) indicates delta-5 desaturase activity.

The negative control strain was *S. cerevisiae* transformed with the pYX242 vector. The experimental and control cultures were grown simultaneously and analyzed.

The cultures were vigorously agitated (250 rpm) and grown for 48 hours at 24°C in the presence of 50 µM (final concentration) of the various substrates (see Table 7). The cells were spun down, washed once in distilled water, and the pellets resuspended in methanol; chloroform was added along with tritridecanoin (as an internal standard). These mixtures were incubated for at least an hour at room temperature, or at 4°C overnight. The chloroform layer was extracted and filtered through a Whatman filter with 1 gm anhydrous sodium sulfate to remove particulate matter and residual water. The organic solvents were evaporated at 40°C under a stream of nitrogen. The extracted lipids were then converted to fatty acid methyl esters (FAME) for gas chromatography analysis (GC) by adding 2 ml 0.5 N potassium hydroxide in methanol to a closed tube. The samples were heated to 95°C-100°C for 30 minutes and cooled to room temperature. Approximately 2 ml 14% borontrifluoride in methanol was added and the heating repeated. After the extracted lipid mixture cooled, 2 ml of water and 1 ml of hexane were added to extract the FAME for analysis by GC. The percent conversion was calculated using the formula:

$$\% \text{ Conversion} = \frac{[\% \text{ Product}]}{[\% \text{ Product} + \% \text{ Substrate}]} \times 100$$

Table 7 shows the enzyme activity of the delta-12 desaturase when expressed in yeast. Here, the pRSP11 clone when expressed was capable of converting 35.8% of OA substrate to LA, indicating delta-12 desaturase activity.

In Table 7, the fatty acids of interest are represented as a percentage of the total lipids extracted from yeast. GC/MS was employed to identify the products. Under these conditions, the clones did not exhibit other desaturase activities. This confirmed that the gene isolated is a delta-12 desaturase gene. No background substrate conversion was detected using hosts transformed with just the vector alone. This data indicates that this delta-12 desaturase can be expressed in a heterologous system and is thus useful in the production of transgenic polyunsaturated fatty acids like GLA, AA, EPA and DHA.

Table 7
***Saprolegnia diclina* Delta 12-Desaturase Expression in Baker's Yeast at 24°C**

Clone	Desatur. Activity	Substrate* Incorporated	Product Produced	% Conversion of Substrate
pRSP11	Delta 12	OA (17.09%)	LA (9.59%)	35.8%
(pYX242 +	Delta 15	LA (18.14%)	ALA (0.06%)	0
Delta 12-	Delta 6	LA (18.14%)	GLA (0%)	0
Desaturase	Delta 5	DGLA (25.38%)	AA (0.17 %)	0
(S.	Delta 17	DGLA (25.38%)	ETA (0.07%)	0
<i>diclina</i>))				
Control	Delta 12	OA (18.99%)	LA (0.09%)	0
(pYX242)	Delta 15	LA (8.63%)	ALA (0%)	0
	Delta 6	LA (8.63%)	GLA (0%)	0
	Delta 5	DGLA (13.74%)	AA (0%)	0
	Delta 17	DGLA (13.74%)	ETA (0%)	0

*50 μ M substrate used

Numbers in parenthesis represent fatty acid as a percentage of total lipids from yeast.

Nutritional Compositions

The PUFAs described in the Detailed Description may be utilized in various nutritional supplements, infant formulations, nutritional substitutes and other nutritional solutions.

I. INFANT FORMULATIONS

A. Isomil® Soy Formula with Iron:

Usage: As a beverage for infants, children and adults with an allergy or sensitivity to cows milk. A feeding for patients with disorders for which lactose should be avoided: lactase deficiency, lactose intolerance and galactosemia.

Features:

- Soy protein isolate to avoid symptoms of cow's-milk-protein allergy or sensitivity.
- Lactose-free formulation to avoid lactose-associated diarrhea.
- Low osmolality (240 mOs/kg water) to reduce risk of osmotic diarrhea.
- Dual carbohydrates (corn syrup and sucrose) designed to enhance carbohydrate absorption and reduce the risk of exceeding the absorptive capacity of the damaged gut.
- 1.8 mg of Iron (as ferrous sulfate) per 100 Calories to help prevent iron deficiency.
- Recommended levels of vitamins and minerals.
- Vegetable oils to provide recommended levels of essential fatty acids.
- Milk-white color, milk-like consistency and pleasant aroma.

Ingredients: (Pareve) 85% water, 4.9% corn syrup, 2.6% sugar (sucrose), 2.1 % soy oil, 1.9% soy protein isolate, 1.4% coconut oil, 0.15% calcium citrate, 0.11 % calcium phosphate tribasic, potassium citrate, potassium phosphate monobasic, potassium chloride, mono- and diglycerides, soy lecithin, carrageenan, ascorbic acid, L-methionine,

magnesium chloride, potassium phosphate dibasic, sodium chloride, choline chloride, taurine, ferrous sulfate, m-inositol, alpha-tocopheryl acetate, zinc sulfate, L-carnitine, niacinamide, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, potassium iodide, phylloquinone, biotin, sodium selenite, vitamin D3 and cyanocobalamin.

B. Isomil® DF Soy Formula For Diarrhea:

Usage: As a short-term feeding for the dietary management of diarrhea in infants and toddlers.

Features:

- First infant formula to contain added dietary fiber from soy fiber specifically for diarrhea management.
- Clinically shown to reduce the duration of loose, watery stools during mild to severe diarrhea in infants.
- Nutritionally complete to meet the nutritional needs of the infant.
- Soy protein isolate with added L-methionine meets or exceeds an infant's requirement for all essential amino acids.
- Lactose-free formulation to avoid lactose-associated diarrhea.
- Low osmolality (240 mOsm/kg water) to reduce the risk of osmotic diarrhea.
- Dual carbohydrates (corn syrup and sucrose) designed to enhance carbohydrate absorption and reduce the risk of exceeding the absorptive capacity of the damaged gut.
- Meets or exceeds the vitamin and mineral levels recommended by the Committee on Nutrition of the American Academy of Pediatrics and required by the Infant Formula Act.

-1.8 mg of iron (as ferrous sulfate) per 100 Calories to help prevent iron deficiency.

-Vegetable oils to provide recommended levels of essential fatty acids.

Ingredients: (Pareve) 86% water, 4.8% corn syrup, 2.5% sugar (sucrose), 2.1% soy oil, 2.0% soy protein isolate, 1.4% coconut oil, 0.77% soy fiber, 0.12% calcium citrate, 0.11% calcium phosphate tribasic, 0.10% potassium citrate, potassium chloride, potassium phosphate monobasic, mono and diglycerides, soy lecithin, carrageenan, magnesium chloride, ascorbic acid, L-methionine, potassium phosphate dibasic, sodium chloride, choline chloride, taurine, ferrous sulfate, m-inositol, alpha-tocopheryl acetate, zinc sulfate, L-carnitine, niacinamide, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, potassium iodide, phylloquinone, biotin, sodium selenite, vitamin D3 and cyanocobalamin.

C. Isomil® SF Sucrose-Free Soy Formula With Iron:

Usage: As a beverage for infants, children and adults with an allergy or sensitivity to cow's-milk protein or an intolerance to sucrose. A feeding for patients with disorders for which lactose and sucrose should be avoided.

Features:

- Soy protein isolate to avoid symptoms of cow's-milk-protein allergy or sensitivity.
- Lactose-free formulation to avoid lactose-associated diarrhea (carbohydrate source is Polycose® Glucose Polymers).
- Sucrose free for the patient who cannot tolerate sucrose.

- Low osmolality (180 mOsm/kg water) to reduce risk of osmotic diarrhea.
- 1.8 mg of iron (as ferrous sulfate) per 100 Calories to help prevent iron deficiency.
- Recommended levels of vitamins and minerals.
- Vegetable oils to provide recommended levels of essential fatty acids.
- Milk-white color, milk-like consistency and pleasant aroma.

Ingredients: (Pareve) 75% water, 11.8% hydrolyzed cornstarch, 4.1% soy oil, 4.1 % soy protein isolate, 2.8% coconut oil, 1.0% modified cornstarch, 0.38% calcium phosphate tribasic, 0. 17% potassium citrate, 0.13% potassium chloride, mono- and diglycerides, soy lecithin, magnesium chloride, ascorbic acid, L-methionine, calcium carbonate, sodium chloride, choline chloride, carrageenan, taurine, ferrous sulfate, m-inositol, alpha-tocopheryl acetate, zinc sulfate, L-carnitine, niacinamide, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, potassium iodide, phylloquinone, biotin, sodium selenite, vitamin D3 and cyanocobalamin.

D. Isomil® 20 Soy Formula With Iron Ready To Feed,
20 Cal/fl oz.:

Usage: When a soy feeding is desired.

Ingredients: (Pareve) 85% water, 4.9% corn syrup, 2.6% sugar(sucrose), 2.1 % soy oil, 1.9% soy protein isolate, 1.4% coconut oil, 0.15% calcium citrate, 0. 11% calcium phosphate tribasic, potassium citrate, potassium phosphate monobasic, potassium chloride, mono- and diglycerides, soy lecithin, carrageenan, ascorbic acid, L-methionine, magnesium chloride, potassium phosphate dibasic, sodium chloride, choline chloride, taurine, ferrous sulfate, m-

inositol, alpha-tocopheryl acetate, zinc sulfate, L-carnitine, niacinamide, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, potassium iodide, phylloquinone, biotin, sodium selenite, vitamin D3 and cyanocobalamin.

E. Similac® Infant Formula:

Usage: When an infant formula is needed: if the decision is made to discontinue breastfeeding before age 1 year, if a supplement to breastfeeding is needed or as a routine feeding if breastfeeding is not adopted.

Features:

- Protein of appropriate quality and quantity for good growth; heat-denatured, which reduces the risk of milk-associated enteric blood loss.
- Fat from a blend of vegetable oils (doubly homogenized), providing essential linoleic acid that is easily absorbed.
- Carbohydrate as lactose in proportion similar to that of human milk.
- Low renal solute load to minimize stress on developing organs.
- Powder, Concentrated Liquid and Ready To Feed forms.

Ingredients: (-D) Water, nonfat milk, lactose, soy oil, coconut oil, mono- and diglycerides, soy lecithin, ascorbic acid, carrageenan, choline chloride, taurine, m-inositol, alpha-tocopheryl acetate, zinc sulfate, niacinamide, ferrous sulfate, calcium pantothenate, cupric sulfate, vitamin A palmitate, thiamine chloride hydrochloride, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, phylloquinone, biotin, sodium selenite, vitamin D3 and cyanocobalamin.

F.Similac® NeoCare Premature Infant Formula With Iron:

Usage: For premature infants' special nutritional needs after hospital discharge. Similac NeoCare is a nutritionally complete formula developed to provide premature infants with extra calories, protein, vitamins and minerals needed to promote catch-up growth and support development.

Features:

- Reduces the need for caloric and vitamin supplementation. More calories (22 Cal/fl oz) than standard term formulas (20 Cal/fl oz).
- Highly absorbed fat blend, with medium-chain triglycerides (MCToil) to help meet the special digestive needs of premature infants.
- Higher levels of protein, vitamins and minerals per 100 calories to extend the nutritional support initiated in-hospital.
- More calcium and phosphorus for improved bone mineralization.

Ingredients: -D Corn syrup solids, nonfat milk, lactose, whey protein concentrate, soy oil, high-oleic safflower oil, fractionated coconut oil (medium chain triglycerides), coconut oil, potassium citrate, calcium phosphate tribasic, calcium carbonate, ascorbic acid, magnesium chloride, potassium chloride, sodium chloride, taurine, ferrous sulfate, m-inositol, choline chloride, ascorbyl palmitate, L-carnitine, alpha-tocopheryl acetate, zinc sulfate, niacinamide, mixed tocopherols, sodium citrate, calcium pantothenate, cupric sulfate, thiamine chloride hydrochloride, vitamin A palmitate, beta carotene, riboflavin, pyridoxine hydrochloride, folic acid, manganese sulfate, phylloquinone, biotin, sodium selenite, vitamin D3 and cyanocobalamin.

G. Similac Natural Care Low-Iron Human Milk Fortifier Ready To Use, 24 Cal/fl oz.:

Usage: Designed to be mixed with human milk or to be fed alternatively with human milk to low-birth-weight infants.

Ingredients: -D Water, nonfat milk, hydrolyzed cornstarch, lactose, fractionated coconut oil (medium-chain triglycerides), whey protein concentrate, soy oil, coconut oil, calcium phosphate tribasic, potassium citrate, magnesium chloride, sodium citrate, ascorbic acid, calcium carbonate, mono and diglycerides, soy lecithin, carrageenan, choline chloride, m-inositol, taurine, niacinamide, L-carnitine, alpha tocopheryl acetate, zinc sulfate, potassium chloride, calcium pantothenate, ferrous sulfate, cupric sulfate, riboflavin, vitamin A palmitate, thiamine chloride hydrochloride, pyridoxine hydrochloride, biotin, folic acid, manganese sulfate, phylloquinone, vitamin D3, sodium selenite and cyanocobalamin.

Various PUFAs of this invention can be substituted and/or added to the infant formulae described above and to other infant formulae known to those in the art.

II. NUTRITIONAL FORMULATIONS

A. ENSURE®

Usage: ENSURE is a low-residue liquid food designed primarily as an oral nutritional supplement to be used with or between meals or, in appropriate amounts, as a meal replacement. ENSURE is lactose- and gluten-free, and is suitable for use in modified diets, including low-cholesterol diets. Although it is primarily an oral supplement, it can be fed by tube.

Patient Conditions:

- For patients on modified diets
- For elderly patients at nutrition risk
- For patients with involuntary weight loss
- For patients recovering from illness or surgery
- For patients who need a low-residue diet

Ingredients: -D Water, Sugar (Sucrose), Maltodextrin (Corn), Calcium and Sodium Caseinates, High-Oleic Safflower Oil, Soy Protein Isolate, Soy Oil, Canola Oil, Potassium Citrate, Calcium Phosphate Tribasic, Sodium Citrate, Magnesium Chloride, Magnesium Phosphate Dibasic, Artificial Flavor, Sodium Chloride, Soy Lecithin, Choline Chloride, Ascorbic Acid, Carrageenan, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Gellan Gum, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Folic Acid, Sodium Molybdate, Chromium Chloride, Biotin, Potassium Iodide, Sodium Selenate.

B.ENSURE® BARS:

Usage: ENSURE BARS are complete, balanced nutrition for supplemental use between or with meals. They provide a delicious, nutrient-rich alternative to other snacks. ENSURE BARS contain <1 g lactose/bar, and Chocolate Fudge Brownie flavor is gluten-free. (Honey Graham Crunch flavor contains gluten.)

Patient Conditions:

- For patients who need extra calories, protein, vitamins and minerals.
- Especially useful for people who do not take in enough calories and nutrients.
- For people who have the ability to chew and swallow

-Not to be used by anyone with a peanut allergy or any type of allergy to nuts.

Ingredients: Honey Graham Crunch -- High-Fructose Corn Syrup, Soy Protein Isolate, Brown Sugar, Honey, Maltodextrin (Corn), Crisp Rice (Milled Rice, Sugar [Sucrose], Salt [Sodium Chloride] and Malt), Oat Bran, Partially Hydrogenated Cottonseed and Soy Oils, Soy Polysaccharide, Glycerine, Whey Protein Concentrate, Polydextrose, Fructose, Calcium Caseinate, Cocoa Powder, Artificial Flavors, Canola Oil, High-Oleic Safflower Oil, Nonfat Dry Milk, Whey Powder, Soy Lecithin and Corn Oil. Manufactured in a facility that processes nuts.

Vitamins and Minerals: Calcium Phosphate Tribasic, Potassium Phosphate Dibasic, Magnesium Oxide, Salt (Sodium Chloride), Potassium Chloride, Ascorbic Acid, Ferric Orthophosphate, Alpha-Tocopheryl Acetate, Niacinamide, Zinc Oxide, Calcium Pantothenate, Copper Gluconate, Manganese Sulfate, Riboflavin, Beta Carotene, Pyridoxine Hydrochloride, Thiamine Mononitrate, Folic Acid, Biotin, Chromium Chloride, Potassium Iodide, Sodium Selenate, Sodium Molybdate, Phylloquinone, Vitamin D3 and Cyanocobalamin.

Protein: Honey Graham Crunch - The protein source is a blend of soy protein isolate and milk proteins.

Soy protein isolate	74%
Milk proteins	26%

Fat: Honey Graham Crunch - The fat source is a blend of partially hydrogenated cottonseed and soybean, canola, high oleic safflower, oils, and soy lecithin.

Partially hydrogenated cottonseed and soybean oil	76%
Canola oil	8%

High-oleic safflower oil	8%
Corn oil	4%
Soy lecithin	4%

Carbohydrate: Honey Graham Crunch - The carbohydrate source is a combination of high-fructose corn syrup, brown sugar, maltodextrin, honey, crisp rice, glycerine, soy polysaccharide, and oat bran.

High-fructose corn syrup	24%
Brown sugar	21%
Maltodextrin	12%
Honey	11%
Crisp rice	9%
Glycerine	9%
Soy Polysaccharide	7%
Oat bran	7%

C. ENSURE® HIGH PROTEIN:

Usage: ENSURE HIGH PROTEIN is a concentrated, high-protein liquid food designed for people who require additional calories, protein, vitamins, and minerals in their diets. It can be used as an oral nutritional supplement with or between meals or, in appropriate amounts, as a meal replacement. ENSURE HIGH PROTEIN is lactose- and gluten-free, and is suitable for use by people recovering from general surgery or hip fractures and by patients at risk for pressure ulcers.

Patient Conditions:

-For patients who require additional calories, protein, vitamins, and minerals, such as patients recovering from general surgery or hip fractures, patients at risk for pressure ulcers, and patients on low-cholesterol diets.

Features:

- Low in saturated fat
- Contains 6 g of total fat and < 5 mg of cholesterol per serving
- Rich, creamy taste
- Excellent source of protein, calcium, and other essential vitamins and minerals
- For low-cholesterol diets
- Lactose-free, easily digested

Ingredients:

Vanilla Supreme: -D Water, Sugar (Sucrose), Maltodextrin (Corn), Calcium and Sodium Caseinates, High-Oleic Safflower Oil, Soy Protein Isolate, Soy Oil, Canola Oil, Potassium Citrate, Calcium Phosphate Tribasic, Sodium Citrate, Magnesium Chloride, Magnesium Phosphate Dibasic, Artificial Flavor, Sodium Chloride, Soy Lecithin, Choline Chloride, Ascorbic Acid, Carrageenan, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Gellan Gum, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Folic Acid, Sodium Molybdate, Chromium Chloride, Biotin, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin D3 and Cyanocobalamin.

Protein:

The protein source is a blend of two high-biologic-value proteins: casein and soy.

Sodium and calcium caseinates 85%

Soy protein isolate 15%

Fat:

The fat source is a blend of three oils: high-oleic safflower, canola, and soy.

High-oleic safflower oil	40%
Canola oil	30%
Soy oil	30%

The level of fat in ENSURE HIGH PROTEIN meets American Heart Association (AHA) guidelines. The 6 grams of fat in ENSURE HIGH PROTEIN represent 24% of the total calories, with 2.6% of the fat being from saturated fatty acids and 7.9% from polyunsaturated fatty acids. These values are within the AHA guidelines of < 30% of total calories from fat, < 10% of the calories from saturated fatty acids, and < 10% of total calories from polyunsaturated fatty acids.

Carbohydrate:

ENSURE HIGH PROTEIN contains a combination of maltodextrin and sucrose. The mild sweetness and flavor variety (vanilla supreme, chocolate royal, wild berry, and banana), plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, help to prevent flavor fatigue and aid in patient compliance.

Vanilla and other nonchocolate flavors:

Sucrose	60%
Maltodextrin	40%

Chocolate:

Sucrose	70%
Maltodextrin	30%

D. ENSURE® LIGHT

Usage: ENSURE LIGHT is a low-fat liquid food designed for use as an oral nutritional supplement with or between meals. ENSURE LIGHT is lactose- and gluten-free, and is suitable for use in modified diets, including low-cholesterol diets.

Patient Conditions:

- For normal-weight or overweight patients who need extra nutrition in a supplement that contains 50% less fat and 20% fewer calories than ENSURE.
- For healthy adults who do not eat right and need extra nutrition.

Features:

- Low in fat and saturated fat
- Contains 3 g of total fat per serving and < 5 mg cholesterol
- Rich, creamy taste
- Excellent source of calcium and other essential vitamins and minerals
- For low-cholesterol diets
- Lactose-free, easily digested

Ingredients:

French Vanilla: -D Water, Maltodextrin (Corn), Sugar (Sucrose), Calcium Caseinate, High-Oleic Safflower Oil, Canola Oil, Magnesium Chloride, Sodium Citrate, Potassium Citrate, Potassium Phosphate Dibasic, Magnesium Phosphate Dibasic, Natural and Artificial Flavor, Calcium Phosphate Tribasic, Cellulose Gel, Choline Chloride, Soy Lecithin, Carrageenan, Salt (Sodium Chloride), Ascorbic Acid, Cellulose Gum, Ferrous Sulfate, Alpha-Tocopheryl Acetate,

Zinc Sulfate, Niacinamide, Manganese Sulfate, Calcium Pantothenate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Vitamin A Palmitate, Pyridoxine Hydrochloride, Riboflavin, Chromium Chloride, Folic Acid, Sodium Molybdate, Biotin, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin D3 and Cyanocobalamin.

Protein:

The protein source is calcium caseinate.

Calcium caseinate	100%
-------------------	------

Fat:

The fat source is a blend of two oils: high-oleic safflower and canola.

High-oleic safflower oil	70%
Canola oil	30%

The level of fat in ENSURE LIGHT meets American Heart Association (AHA) guidelines. The 3 grams of fat in ENSURE LIGHT represent 13.5% of the total calories, with 1.4% of the fat being from saturated fatty acids and 2.6% from polyunsaturated fatty acids. These values are within the AHA guidelines of < 30% of total calories from fat, < 10% of the, calories from saturated fatty acids, and < 10% of total calories from polyunsaturated fatty acids.

Carbohydrate:

ENSURE LIGHT contains a combination of maltodextrin and sucrose. The chocolate flavor contains corn syrup as well. The mild sweetness and flavor variety (French vanilla, chocolate supreme, strawberry swirl), plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and

orange, help to prevent flavor fatigue and aid in patient compliance.

Vanilla and other nonchocolate flavors:

Sucrose	51%
Maltodextrin	49%

Chocolate:

Sucrose	47.0%
Corn Syrup	26.5%
Maltodextrin	26.5%

Vitamins and Minerals:

An 8-fl-oz serving of ENSURE LIGHT provides at least 25% of the RDIs for 24 key vitamins and minerals.

Caffeine:

Chocolate flavor contains 2.1 mg caffeine/8 fl oz.

E.ENSURE PLUS®

Usage: ENSURE PLUS is a high-calorie, low-residue liquid food for use when extra calories and nutrients, but a normal concentration of protein, are needed. It is designed primarily as an oral nutritional supplement to be used with or between meals or, in appropriate amounts, as a meal replacement. ENSURE PLUS is lactose- and gluten-free. Although it is primarily an oral nutritional supplement, it can be fed by tube.

Patient Conditions:

- For patients who require extra calories and nutrients, but a normal concentration of protein, in a limited volume.
- For patients who need to gain or maintain healthy weight.

Features:

- Rich, creamy taste
- Good source of essential vitamins and minerals

Ingredients:

Vanilla: -D Water, Corn Syrup, Maltodextrin (Corn), Corn Oil, Sodium and Calcium Caseinates, Sugar (Sucrose), Soy Protein Isolate, Magnesium Chloride, Potassium Citrate, Calcium Phosphate Tribasic, Soy Lecithin, Natural and Artificial Flavor, Sodium Citrate, Potassium Chloride, Choline Chloride, Ascorbic Acid, Carrageenan, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenite, Phylloquinone, Cyanocobalamin and Vitamin D3.

Protein:

The protein source is a blend of two high-biologic-value proteins: casein and soy.

Sodium and calcium caseinates	84%
Soy protein isolate	16%

Fat:

The fat source is corn oil.

Corn oil	100%
----------	------

Carbohydrate:

ENSURE PLUS contains a combination of maltodextrin and sucrose. The mild sweetness and flavor variety (vanilla,

chocolate, strawberry, coffee, butter pecan, and eggnog), plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, help to prevent flavor fatigue and aid in patient compliance.

Vanilla, strawberry, butter pecan, and coffee flavors:

Corn Syrup	39%
Maltodextrin	38%
Sucrose	23%

Chocolate and eggnog flavors:

Corn Syrup	36%
Maltodextrin	34%
Sucrose	30%

Vitamins and Minerals:

An 8-fl-oz serving of ENSURE PLUS provides at least 15% of the RDIs for 25 key Vitamins and minerals.

Caffeine:

Chocolate flavor contains 3.1 mg Caffeine/8 fl oz. Coffee flavor contains a trace amount of caffeine.

F. ENSURE PLUS® HN

Usage: ENSURE PLUS HN is a nutritionally complete high-calorie, high-nitrogen liquid food designed for people with higher calorie and protein needs or limited volume tolerance. It may be used for oral supplementation or for total nutritional support by tube. ENSURE PLUS HN is lactose- and gluten-free.

Patient Conditions:

- For patients with increased calorie and protein needs, such as following surgery or injury.
- For patients with limited volume tolerance and early satiety.

Features:

- For supplemental or total nutrition
- For oral or tube feeding
- 1.5 CaV/mL,
- High nitrogen
- Calorically dense

Ingredients:

Vanilla: -D Water, Maltodextrin (Corn), Sodium and Calcium Caseinates, Corn Oil, Sugar (Sucrose), Soy Protein Isolate, Magnesium Chloride, Potassium Citrate, Calcium Phosphate Tribasic, Soy Lecithin, Natural and Artificial Flavor, Sodium Citrate, Choline Chloride, Ascorbic Acid, Taurine, L-Carnitine, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Niacinamide, Carrageenan, Calcium Pantothenate, Manganese Sulfate, Cupric Sulfate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Folic Acid, Biotin, Chromium Chloride, Sodium Molybdate, Potassium Iodide, Sodium Selenite, Phylloquinone, Cyanocobalamin and Vitamin D3.

G.ENSURE® POWDER:

Usage: ENSURE POWDER (reconstituted with water) is a low-residue liquid food designed primarily as an oral nutritional supplement to be used with or between meals. ENSURE POWDER is lactose- and gluten-free, and is suitable for use in modified diets, including low-cholesterol diets.

Patient Conditions:

- For patients on modified diets
- For elderly patients at nutrition risk
- For patients recovering from illness/surgery
- For patients who need a low-residue diet

Features:

- Convenient, easy to mix
- Low in saturated fat
- Contains 9 g of total fat and < 5 mg of cholesterol per serving
- High in vitamins and minerals
- For low-cholesterol diets
- Lactose-free, easily digested

Ingredients: -D Corn Syrup, Maltodextrin (Corn), Sugar (Sucrose), Corn Oil, Sodium and Calcium Caseinates, Soy Protein Isolate, Artificial Flavor, Potassium Citrate, Magnesium Chloride, Sodium Citrate, Calcium Phosphate Tribasic, Potassium Chloride, Soy Lecithin, Ascorbic Acid, Choline Chloride, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Niacinamide, Calcium Pantothenate, Manganese Sulfate, Thiamine Chloride Hydrochloride, Cupric Sulfate, Pyridoxine Hydrochloride, Riboflavin, Vitamin A Palmitate, Folic Acid, Biotin, Sodium Molybdate, Chromium Chloride, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin D3 and Cyanocobalamin.

Protein:

The protein source is a blend of two high-biologic-value proteins: casein and soy.

Sodium and calcium caseinates	84%
Soy protein isolate	16%

Fat:

The fat source is corn oil.

Corn oil

100%

Carbohydrate:

ENSURE POWDER contains a combination of corn syrup, maltodextrin, and sucrose. The mild sweetness of ENSURE POWDER, plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, helps to prevent flavor fatigue and aid in patient compliance.

Vanilla:

Corn Syrup

35%

Maltodextrin

35%

Sucrose

30%

H. ENSURE® PUDDING

Usage: ENSURE PUDDING is a nutrient-dense supplement providing balanced nutrition in a nonliquid form to be used with or between meals. It is appropriate for consistency-modified diets (e.g., soft, pureed, or full liquid) or for people with swallowing impairments. ENSURE PUDDING is gluten-free.

Patient Conditions:

- For patients on consistency-modified diets (e.g., soft, pureed, or full liquid)
- For patients with swallowing impairments

Features:

- Rich and creamy, good taste
- Good source of essential vitamins and minerals

- Convenient-needs no refrigeration
- Gluten-free

Nutrient Profile per 5 oz: Calories 250, Protein 10.9%,
Total Fat 34.9%, Carbohydrate 54.2%

Ingredients:

Vanilla: -D Nonfat Milk, Water, Sugar (Sucrose), Partially Hydrogenated Soybean Oil, Modified Food Starch, Magnesium Sulfate, Sodium Stearoyl Lactylate, Sodium Phosphate Dibasic, Artificial Flavor, Ascorbic Acid, Zinc Sulfate, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Choline Chloride, Niacinamide, Manganese Sulfate, Calcium Pantothenate, FD&C Yellow #5, Potassium Citrate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, FD&C Yellow #6, Folic Acid, Biotin, Phylloquinone, Vitamin D3 and Cyanocobalamin.

Protein:

The protein source is nonfat milk.

Nonfat milk	100%
-------------	------

Fat:

The fat source is hydrogenated soybean oil.

Hydrogenated soybean oil	100%
--------------------------	------

Carbohydrate:

ENSURE PUDDING contains a combination of sucrose and modified food starch. The mild sweetness and flavor variety (vanilla, chocolate, butterscotch, and tapioca) help prevent flavor fatigue. The product contains 9.2 grams of lactose per serving.

Vanilla and other nonchocolate flavors:

Sucrose	56%
Lactose	27%
Modified food starch	17%

Chocolate:

Sucrose	58%
Lactose	26%
Modified food starch	16%

I. ENSURE® WITH FIBER:

Usage: ENSURE WITH FIBER is a fiber-containing, nutritionally complete liquid food designed for people who can benefit from increased dietary fiber and nutrients. ENSURE WITH FIBER is suitable for people who do not require a low-residue diet. It can be fed orally or by tube, and can be used as a nutritional supplement to a regular diet or, in appropriate amounts, as a meal replacement. ENSURE WITH FIBER is lactose- and gluten-free, and is suitable for use in modified diets, including low-cholesterol diets.

Patient Conditions:

-For patients who can benefit from increased dietary fiber and nutrients

Features:

- New advanced formula-low in saturated fat, higher in vitamins and minerals
- Contains 6 g of total fat and < 5 mg of cholesterol per serving
- Rich, creamy taste
- Good source of fiber

- Excellent source of essential vitamins and minerals
- For low-cholesterol diets
- Lactose- and gluten-free

Ingredients:

Vanilla: -D Water; Maltodextrin (Corn), Sugar (Sucrose), Sodium and Calcium Caseinates, Oat Fiber, High-Oleic Safflower Oil, Canola Oil, Soy Protein Isolate, Corn Oil, Soy Fiber, Calcium Phosphate Tribasic, Magnesium Chloride, Potassium Citrate, Cellulose Gel, Soy Lecithin, Potassium Phosphate Dibasic, Sodium Citrate, Natural and Artificial Flavors, Choline Chloride, Magnesium Phosphate, Ascorbic Acid, Cellulose Gum, Potassium Chloride, Carrageenan, Ferrous Sulfate, Alpha-Tocopheryl Acetate, Zinc Sulfate, Niacinamide, Manganese Sulfate, Calcium Pantothenate, Cupric Sulfate, Vitamin A Palmitate, Thiamine Chloride Hydrochloride, Pyridoxine Hydrochloride, Riboflavin, Folic Acid, Chromium Chloride, Biotin, Sodium Molybdate, Potassium Iodide, Sodium Selenate, Phylloquinone, Vitamin D3 and Cyanocobalamin.

Protein:

The protein source is a blend of two high-biologic-value proteins-casein and soy.

Sodium and calcium caseinates	80%
Soy protein isolate	20%

Fat:

The fat source is a blend of three oils: high-oleic safflower, canola, and corn.

High-oleic safflower oil	40%
Canola oil	40%
Corn oil	20%

The level of fat in ENSURE WITH FIBER meets American Heart Association (AHA) guidelines. The 6 grams of fat in ENSURE WITH FIBER represent 22% of the total calories, with 2.01 % of the fat being from saturated fatty acids and 6.7% from polyunsaturated fatty acids. These values are within the AHA guidelines of $\leq 30\%$ of total calories from fat, $< 10\%$ of the calories from saturated fatty acids, and $\leq 10\%$ of total calories from polyunsaturated fatty acids.

Carbohydrate:

ENSURE WITH FIBER contains a combination of maltodextrin and sucrose. The mild sweetness and flavor variety (vanilla, chocolate, and butter pecan), plus VARI-FLAVORS® Flavor Pacs in pecan, cherry, strawberry, lemon, and orange, help to prevent flavor fatigue and aid in patient compliance.

Vanilla and other nonchocolate flavors:

Maltodextrin	66%
Sucrose	25%
Oat Fiber	7%
Soy Fiber	2%

Chocolate:

Maltodextrin	55%
Sucrose	36%
Oat Fiber	7%
Soy Fiber	2%

Fiber:

The fiber blend used in ENSURE WITH FIBER consists of oat fiber and soy polysaccharide. This blend results in

approximately 4 grams of total dietary fiber per 8-fl. oz can. The ratio of insoluble to soluble fiber is 95:5.

The various nutritional supplements described above and known to others of skill in the art can be substituted and/or supplemented with the PUFAs produced in accordance with the present invention.

J.Oxepa™ Nutritional Product

Oxepa is a low-carbohydrate, calorically dense, enteral nutritional product designed for the dietary management of patients with or at risk for ARDS. It has a unique combination of ingredients, including a patented oil blend containing eicosapentaenoic acid (EPA from fish oil), γ -linolenic acid (GLA from borage oil), and elevated antioxidant levels.

Caloric Distribution:

Caloric density is high at 1.5 Cal/mL (355 Cal/8 fl oz), to minimize the volume required to meet energy needs.

The distribution of Calories in Oxepa is shown in Table A.

Table A. Caloric Distribution of Oxepa

	per 8 fl oz.	per liter	% of Cal
Calories	355	1,500	---
Fat (g)	22.2	93.7	55.2
Carbohydrate (g)		25	105.528.1
Protein (g)	14.8	62.5	16.7
Water (g)	186	785	---

Fat:

-Oxepa contains 22.2 g of fat per 8-fl oz serving (93.7 g/L).

-The fat source is an oil blend of 31.8% canola oil, 25% medium-chain triglycerides (MCTs), 20% borage oil, 20% fish

oil, and 3.2 % soy lecithin. The typical fatty acid profile of Oxepa is shown in Table B.

-Oxepa provides a balanced amount of polyunsaturated, monounsaturated, and saturated fatty acids, as shown in Table VI.

-Medium-chain triglycerides (MCTs) -- 25% of the fat blend -
- aid gastric emptying because they are absorbed by the intestinal tract without emulsification by bile acids.

The various fatty acid components of Oxepa™ nutritional product can be substituted and/or supplemented with the PUFAs produced in accordance with this invention.

Table B. Typical Fatty Acid Profile

	% Total	g/8 fl oz*	g/L*
Fatty Acids			
Caproic (6:0)	0.2	0.04	0.18
Caprylic (8:0)	14.69	3.1	13.07
Capric (10:0)	11.06	2.33	9.87
Palmitic (16:0)	5.59	1.18	4.98
Palmitoleic	1.82	0.38	1.62
Stearic	1.94	0.39	1.64
Oleic	24.44	5.16	21.75
Linoleic	16.28	3.44	14.49
α -Linolenic	3.47	0.73	3.09
γ -Linolenic	4.82	1.02	4.29
Eicosapentaenoic	5.11	1.08	4.55
n-3-Docosapent- aenoic	0.55	0.12	0.49
Docosaheptaenoic	2.27	0.48	2.02

Others	7.55	1.52	6.72
--------	------	------	------

Fatty acids equal approximately 95% of total fat.

Table C. Fat Profile of Oxepa

% of total calories from fat	55.2
Polyunsaturated fatty acids	31.44 g/L
Monounsaturated fatty acids	25.53 g/L
Saturated fatty acids	32.38 g/L
n-6 to n-3 ratio	1.75:1
Cholesterol	9.49 mg/8 fl oz
	40.1 mg/L

Carbohydrate:

-The carbohydrate content is 25.0 g per 8-fl-oz serving (105.5 g/L).

-The carbohydrate sources are 45% maltodextrin (a complex carbohydrate) and 55% sucrose (a simple sugar), both of which are readily digested and absorbed.

-The high-fat and low-carbohydrate content of Oxepa is designed to minimize carbon dioxide (CO₂) production. High CO₂ levels can complicate weaning in ventilator-dependent patients. The low level of carbohydrate also may be useful for those patients who have developed stress-induced hyperglycemia.

-Oxepa is lactose-free.

Dietary carbohydrate, the amino acids from protein, and the glycerol moiety of fats can be converted to glucose within the body. Throughout this process, the carbohydrate requirements of glucose-dependent tissues (such as the central nervous system and red blood cells) are met. However, a diet free of carbohydrates can lead to ketosis, excessive catabolism of tissue protein, and loss of fluid and electrolytes. These effects can be prevented by daily

ingestion of 50 to 100 g of digestible carbohydrate, if caloric intake is adequate. The carbohydrate level in Oxepa is also sufficient to minimize gluconeogenesis, if energy needs are being met.

Protein:

-Oxepa contains 14.8 g of protein per 8-fl-oz serving (62.5 g/L).

-The total calorie/nitrogen ratio (150:1) meets the need of stressed patients.

-Oxepa provides enough protein to promote anabolism and the maintenance of lean body mass without precipitating respiratory problems. High protein intakes are a concern in patients with respiratory insufficiency. Although protein has little effect on CO₂ production, a high protein diet will increase ventilatory drive.

-The protein sources of Oxepa are 86.8% sodium caseinate and 13.2% calcium caseinate.

- The amino acid profile of the protein system in Oxepa meets or surpasses the standard for high quality protein set by the National Academy of Sciences.

* Oxepa is gluten-free.

CLAIMS

What is claimed is:

1. An isolated nucleotide acid sequence or fragment thereof comprising or complementary to a nucleotide sequence encoding a polypeptide having desaturase activity, wherein the amino acid sequence of said polypeptide has at least 50% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:26 and SEQ ID NO:42.

2. An isolated nucleotide sequence or fragment thereof comprising or complementary to at least 50% of the nucleotide sequence selected from the group consisting of SEQ ID NO:25 and SEQ ID NO:41.

3. The isolated nucleotide sequence of claim 2, wherein said sequence is selected from the group consisting of SEQ ID NO:25 and SEQ ID NO:41.

4. The isolated nucleotide sequence of claim 2 or 3, wherein said sequence encodes a functionally active desaturase which utilizes a polyunsaturated fatty acid as a substrate.

5. The isolated nucleotide sequence of claim 1 or 2, wherein said sequence is derived from *Saprolegnia diclina*.

6. A purified polypeptide encoded by said isolated nucleotide sequence of claim 1 or 2.

7. A purified polypeptide that desaturates a polyunsaturated fatty acid substrate at an omega-3 carbon of said substrate and has at least 50% amino acid identity to an amino acid sequence comprising SEQ ID NO: 26.

8. The purified polypeptide of claim 7, wherein said

polypeptide desaturates a fatty acid substrate having 20 carbon atoms.

9. A purified polypeptide that desaturates a polyunsaturated fatty acid substrate at a delta-12 carbon of said substrate and has at least 50% amino acid identity to SEQ ID NO: 42.

10. The purified polypeptide of claim 9, wherein said polypeptide desaturates a fatty acid substrate having 18 carbon atoms.

11. A method of producing a desaturase comprising the steps of:

- (a) isolating a nucleotide sequence comprising or complementary to at least 50% of the nucleotide sequence selected from the group consisting of SEQ ID NO: 25 and SEQ ID NO: 41;
- (b) constructing a vector comprising said isolated nucleotide sequence of step (a); and
- (c) introducing said vector of step (b) into a host cell for a time and under conditions sufficient for expression of a desaturase encoded by said isolated nucleotide sequence of step (a).

12. A vector comprising: 1) an isolated nucleotide sequence corresponding to or complementary to at least 50% of the nucleotide sequence selected from the group consisting of SEQ ID NO: 25 and SEQ ID NO: 41, operably linked to b) a regulatory sequence.

13. A host cell comprising said vector of claim 12.

14. The host cell of claim 13, wherein said host cell is a eukaryotic cell selected from the group consisting of a mammalian cell, an insect cell, a plant cell and a fungal cell.

15. The host cell of claim 14, wherein expression of said isolated nucleotide sequence of said vector results in said host cell producing a polyunsaturated fatty acid that is not produced in a wild-type of said host cell.

16. A plant cell, plant, or plant tissue comprising said vector of claim 12, wherein expression of said nucleotide sequence of said vector results in production of a polyunsaturated fatty acid by said plant cell, plant or plant tissue.

17. The plant cell, plant, or plant tissue of claim 16, wherein said vector induces the production of a polyunsaturated fatty acid selected from the group consisting of linoleic acid, eicosatetraenoic acid and eicosapentaenoic acid.

18. A transgenic plant comprising said vector of claim 12, wherein expression of said nucleotide sequence of said vector results in production of a polyunsaturated fatty acid in seeds of said transgenic plant.

19. One or more plant oils or acids expressed by said plant cell, plant or plant tissue of claim 16.

20. A method for producing a polyunsaturated fatty acid comprising the steps of:

- (a) isolating a nucleotide sequence comprising or complementary to at least 50% of the nucleotide sequence selected from the group consisting of SEQ ID NO: 25 and SEQ ID NO: 41;
- (b) constructing a vector comprising said isolated nucleotide sequence of step (a);
- (c) transforming the vector of step (b) into a host cell for a time and under conditions sufficient for expression of a desaturase encoded by said isolated nucleotide sequence of step (a); and

(d) exposing said expressed desaturase selected from the group consisting of an omega-3-desaturase and a delta 12-desaturase, to a fatty acid substrate, whereby said substrate is catalytically converted by said desaturase into a desired polyunsaturated fatty acid product.

21. The method of claim 20, wherein said substrate is dihomogamma-linolenic acid or arachidonic acid and said product polyunsaturated fatty acid is eicosatetraenoic acid or eicosapentaenoic acid, respectively, when said expressed desaturase is an omega-3-desaturase.

22. The method of claim 20, wherein said substrate polyunsaturated fatty acid is oleic acid and said product polyunsaturated fatty acid is linoleic acid, when said expressed desaturase is a delta 12-desaturase.

23. The method of claim 20, further comprising, after step (d), the step of:
(e) exposing said polyunsaturated fatty acid product of step (d) to one or more enzymes selected from the group consisting of a desaturase and an elongase, whereby the polyunsaturated fatty acid product of step (d) is catalytically converted into another polyunsaturated fatty acid product.

24. The method of claim 23, wherein said product polyunsaturated fatty acid is eicosatetraenoic acid or eicosapentaenoic acid and said another polyunsaturated fatty acid is eicosapentaenoic acid or omega 3-docosapentaenoic acid, respectively, when said expressed desaturase of step (d) is an omega 3-desaturase.

25. The method of claim 23, wherein said product polyunsaturated fatty acid is linoleic acid and said another polyunsaturated fatty acid is gamma-linolenic acid, when said expressed desaturase of step (d) is a delta 12-desaturase.

26. The method of claim 23 further comprising the step of exposing said another polyunsaturated fatty acid to one or more enzymes selected from the group consisting of a desaturase and an elongase in order to convert said another polyunsaturated fatty acid to a final polyunsaturated fatty acid.

27. The method of claim 26 wherein said final polyunsaturated fatty acid is selected from the group consisting of omega 3-docosapentaenoic acid and docosahexaenoic acid, when said expressed desaturase of step (d) is an omega 3-desaturase.

28. The method of claim 26 wherein said final polyunsaturated fatty acid is selected from the group consisting of dihomo-gamma-linolenic acid, arachidonic acid, adrenic acid, omega 6-docosapentaenoic acid, eicosatetraenoic acid, stearidonic acid, eicosapentaenoic acid, omega 3-docosapentaenoic acid and docosahexaenoic acid, when said expressed desaturase of step (d) is a delta 12-desaturase.

29. A method of producing a polyunsaturated fatty acid comprising exposing a fatty acid substrate to a polypeptide having at least 50% amino acid identity to an amino acid sequence selected from the group consisting of SEQ ID NO: 26 and SEQ ID NO: 42, whereby said fatty acid substrate is catalytically converted into said polyunsaturated fatty acid.

30. The method of claim 29, wherein the fatty acid substrate is dihomo-gamma-linolenic acid or arachidonic acid and said product polyunsaturated fatty acid is eicosatetraenoic acid or eicosapentaenoic acid, respectively, when said polypeptide is an omega 3-desaturase.

31. The method of claim 29, wherein said fatty acid substrate is oleic acid and said polyunsaturated fatty acid is linoleic acid, when said polypeptide is a delta 12-desaturase.

32. A composition comprising at least one polyunsaturated fatty acid selected from the group consisting of said product polyunsaturated fatty acid produced according to the method of claim 20, said another polyunsaturated fatty acid produced according to the method of claim 23, and said final polyunsaturated fatty acid produced according to the method of claim 26.

33. The composition of claim 32, wherein said product polyunsaturated fatty acid is eicosatetraenoic acid or eicosapentaenoic acid, when said expressed desaturase of step (d) is an omega 3-desaturase.

34. The composition of claim 32, wherein said product polyunsaturated fatty acid is linoleic acid, when said expressed desaturase of step (d) is a delta 12-desaturase.

35. The composition of claim 32, wherein said another polyunsaturated fatty acid is eicosapentaenoic acid or omega 3-docosapentaenoic acid, respectively, when said expressed desaturase of step (d) is an omega 3-desaturase.

36. The composition of claim 32, wherein said another polyunsaturated fatty acid is gamma-linolenic acid, when said expressed desaturase of step (d) is a delta 12-desaturase.

37. The composition of claim 32, wherein said final polyunsaturated fatty acid is selected from the group consisting of omega 3-docosapentaenoic acid and docosahexaenoic acid, when said expressed desaturase of step (d) is an omega 3-desaturase.

38. The composition of claim 32, wherein said final polyunsaturated fatty acid is selected from the group consisting of dihomo-gamma-linolenic acid, arachidonic acid, adrenic acid, omega 6-docosapentaenoic acid, eicosatetraenoic acid, stearidonic

acid, eicosapentaenoic acid, omega 3-docosapentaenoic acid and docosahexaenoic acid, when said expressed desaturase of step (d) is a delta 12-desaturase.

39. A method of preventing or treating a condition caused by insufficient intake of at least one polyunsaturated fatty acid comprising administering to said patient said composition of claim 32 in an amount sufficient to effect said prevention or treatment.




FIG.1

2/14

Gene Sequence of *sdd17*, an Omega-3 Fatty Acid Desaturase Gene
from *Saprolegnia diclina* (ATCC 56851)

```

1  ATGACTGAGG ATAAGACGAA GGTCGAGTTC CCGACGCTCA CGGAGCTCAA
51  GCACTCGATC CCGAACGCGT GCTTTGAGTC GAACCTCGGC CTCTCGCTCT
101 ACTACACGGC CCGCGCGATC TTCAACGCGT CGGCCTCGGC GGCGCTGCTC
151 TACGCGGCGC GCTCGACGCC GTTCATTGCC GATAACGTTT TGCTCCACGC
201 GCTCGTTTGC GCCACCTACA TCTACGTGCA GGGCGTCATC TTCTGGGGCT
251 TCTTCACGGT CGGCCACGAC TCGGGCCACT CGGCCTTCTC GCGCTACCAC
301 AGCGTCAACT TTATCATCGG CTGCATCATG CACTCTGCGA TTTTGACGCC
351 GTTCGAGAGC TGGCGCGTGA CGCACCGCCA CCACCACAAG AACACGGGCA
401 ACATTGATAA GGACGAGATC TTTTACCCGC ACCGGTCGGT CAAGGACCTC
451 CAGGACGTGC GCCAATGGGT CTACACGCTC GGCGGTGCGT GGTTTGTCTA
501 CTTGAAGGTC GGGTATGCCC CGCGCACGAT GAGCCACTTT GACCCGTGGG
551 ACCCGCTCCT CCTTCGCCGC GCGTCGGCCG TCATCGTGTC GCTCGGCGTC
601 TGGGCCGCCT TCTTCGCCGC GTACGCGTAC CTCACATACT CGCTCGGCTT
651 TGCCGTCATG GGCCTCTACT ACTATGCGCC GCTCTTTGTC TTTGCTTCGT
701 TCCTCGTCAT TACGACCTTC TTGCACCACA ACGACGAAGC GACGCCGTGG
751 TACGGCGACT CGGAGTGGAC GTACGTCAAG GGCAACCTCT CGAGCGTCGA
801 CCGCTCGTAC GGC CGTTCG TGGACAACCT GAGCCACCAC ATTGGCACGC
851 ACCAGGTCCA CCACTTGTTT CCGATCATTG CGCACTACAA GCTCAACGAA
901 GCCACCAAGC ACTTTGCGGC CGCGTACCCG CACCTCGTGC GCAGGAACGA
951 CGAGCCCATC ATCACGGCCT TCTTCAAGAC CGCGCACCTC TTTGTCAACT
1001 ACGGCGCTGT GCCCGAGACG GCGCAGATCT TCACGCTCAA AGAGTCGGCC
1051 GCGGCCGCCA AGGCCAAGTC GGACTAA

```

FIG.2

3/14

Amino Acid Sequence of an Omega-3 Fatty Acid Desaturase (SDD17) from
Saprolegnia diclina (ATCC 56851)

```
1  MTEDKTKVEF PTLTELKHSI PNACFESNLG LSLYYTARAI FNASASAALL
51  YAARSTPFIA DNVLLHALVC ATYIYVQGVI FWGFFTVGHD CGHSAFSRYH
101 SVNFIIGCIM HSAILTPFES WRVTHRHHHK NTGNIDKDEI FYPHRSVKDL
151 QDVRQWVYTL GGAWFVYLKV GYAPRTMSHF DPWDPLLLRR ASAVIVSLGV
201 WAAFFAAYAY LTYSLGFAVM GLYYYAPLFV FASFLVITTF LHHNDEATPW
251 YGDSEWTVVK GNLSSVDRSY GAFVDNLSSH IGTHQVHHLF PIIPHYKLNE
301 ATKHFAAAYP HLVRRNDEPI ITAFFKTAHL FVNYGAVPET AQIFTLKESA
351 AAAKAKSD*
```

FIG.3

Comparative analysis of *S. diclina* Delta 17-desaturase (SDD17.pep)
& *Synechocystis* sp. Delta 15-desaturase (SYCDESB)

	50	60	70	80	90	100
SDD17.pep	SAALLYAARSTPFIADNVLLHALVCATYIYVQGVIFWGFFT	VGHDCGHS	AFSRYHSVNF	I		
				:	:	:
SYCDESB	YFFLDVGLIAGFYALAA	YLD	SWFFYP	IFWL	IQGTLFWSL	FVVGHDCGHSFSKSKTLNNW
	530	560	590	620	650	680
	110	120	130	140	150	160
SDD17.pep	IGCIMHSAILTPFESWRV	THRHHH	KNTGNIDK	DEIFYPHRSVKDLQ	----	DVRQWVYTLG
	:	:	:	:	:	:
SYCDESB	IGHLSHTPILVPYHGW	RISHRTH	HANTGNID	TDES	WYPVSEQKYNQMAWY	EKLLRFYLP
	710	740	770	800	830	860
	170	180	190	200	210	220
SDD17.pep	GAWFVYLKVG	YAPRTMSH	FD	PWDPL	LL	-RRASAVIVSLGWAAFFAAYAL
	:	:	:	:	:	:
SYCDESB	IAYPIYLFRRSP	NRQGS	HFM	PGSPLFR	PG	EKA
	890	920	950	980	1010	1040
	230	240	250	260	270	280
SDD17.pep	GLYYYYAPLFV	FASFLV	ITTF	LHHNDEAT	PWYGDSEW	TYVKG
	:	:	:	:	:	:
SYCDESB	LKFYVAPYLV	FVWLDL	VTFL	HTEDNIP	WYRGDD	WYFLKGALSTIDRDY
	1070	1100	1130	1160	1190	1220
	290	300	310	320	330	
SDD17.pep	IGTHQVHH	LFPII	PHYKLNEAT	KHFAAAYPHL	VRRNDEPIITAFFKT	---AHLFVNYGAV
	:	:	:	:	:	:
SYCDESB	IGTHVAHH	IFSNM	PHYKLRR	ATEAKPIL	GEYYR	SDEPIWQAF
	1250	1280	1310	1340	1370	1400
	340	350				
SDD17.pep	PETAQIF	TLKESAAA	AKAKSD			
SYCDESB	VYYQSPSNGGYQ	KKPXL	ILIESNQ	HREG	RQYX	MVLLPSDRLMRSME
	1430	1460	1490	1520	1550	1580

FIG. 4

Comparative analysis of *S. diclina* Delta 17-desaturase (SDD17.pep)
C. elegans Delta 17-desaturase (CELEFAT)

[illegible]

FIG. 5

6/14

Gene Sequence of *sdd12*, a Delta 12-Desaturase Gene
from *Saprolegnia diclina* (ATCC 56851)

```

1  ATGTGCAAAG GTCAAGCTCC TTCCAAGGCC GACGTGTTCC ACGCTGCGGG
51  GTACCGCCCG GTCGCCGGCA CGCCCGAGCC GCTGCCGCTG GAGCCCCCGA
101 CGATCACGCT CAAGGACCTG CGCGCGGCGA TCCCGGCCCA CTGCTTTGAG
151 CGCAGCGCTG CCACTAGCTT TTACCATTTG GCCAAGAACC TTGCGATCTG
201 CGCCGGCGTG TTCGCCGTTG GCCTCAAGCT CGCGGCTGCC GACTTGCCGC
251 TCGCGGCCAA GCTGGTCGCG TGGCCCATCT ACTGGTTCGT CCAGGGCAGC
301 TACTTTACGG GCATCTGGGT CATTGCGCAC GAATGCGGCC ACCAGGCGTT
351 CTCGGCGTCC GAGATCCTCA ACGACACGGT CGGTATCATT CTTCACTCGC
401 TCCTCTTTGT GCCGTACCAC AGCTGGAAGA TCACGCACCG CCGCCACCAC
451 TCCAACACGG GCAGCTGCGA GAACGACGAG GTGTTTACGC CGACGCCGCG
501 GTCCGTCGTC GAGGCCAAGC ACGACCACTC GCTCCTCGAA GAGAGCCCGC
551 TCTACAACCT GTACGGCATC GTCATGATGC TTCTCGTGGG CTGGATGCCG
601 GGCTACCTCT TCTTCAACGC GACCGGCCCG ACCAAGTACG CTGGCCTCGC
651 CAAGTCGCAC TTCAACCCGT ACGCAGCCTT TTTCTCCCA AAGGAGCGCC
701 TCAGCATCTG GTGGAGCGAC CTCTGCTTCC TCGCGGCCTT GTACGGCTTT
751 GGCTACGGCG TCTCGGTCTT CGGCCTCCTC GATGTCGCCC GCCACTACAT
801 CGTGCCGTAC CTCATTTGCA ACGCGTACCT CGTGCTCATC ACGTACCTCC
851 AGCACACGGA TACGTACGTG CCCCACCTTC GCGGCGACGA GTGGAAGTGG
901 CTGCGCGGCG CGCTCTGCAC CGTCGACCGC TCGTTCGGCG CGTGGATCGA
951 CAGCGCGATC CACCACATTG CCGACACGCA CGTGACGCAC CACATTTTCT
1001 CCAAGACGCC CTTCTACCAC GCGATCGAGG CGACCGACGC CATCACGCCC
1051 CTCCTCGGCA AGTACTACCT CATCGACCCG ACGCCGATCC CGCTGGCGCT
1101 CTGGCGCTCG TTCACGCACT GCAAGTACGT CGAGGACGAC GGCAACGTTG
1151 TGTTTTACAA GCGCAAGCTC GAGGAAAAGT AA

```

FIG.6

7/14

Amino Acid Sequence of α Delta 12-Desaturase (SDD12) from *Saprolegnia diclina* (ATCC 56851)

```
1  MCKGQAPSKA DVFHAAGYRP VAGTPEPLPL EPPTITLKDL RAAIPAHCFE
51  RSAATSFYHL AKNLAICAGV FAVGLKLAAA DLPLAAKLVA WPIYWVQGT
101 YFTGIWVIAH ECGHQAFSAS EILNDTVGII LHSLLFVPYH SWKITHRRHH
151 SNTGSCENDE VFTPTPRSVV EAKHDHSLLE ESPLYNLYGI VMMLLVGWMP
201 GYLFFNATGP TKYAGLAKSH FNPYAAFFLP KERLSIWWSL LCFLAALYGF
251 GYGVSFVGLL DVARHYIVPY LICNAYLVLI TYLQHTDTYV PHFRGDEWNW
301 LRGALCTVDR SFGAWIDSAI HHIADTHVTH HIFSKTPFYH AIEATDAITP
351 LLGKYYLIDP TPIPLALWRS FTHCKYVEDD GNVVFYKRKL EEK*
```

FIG.7

8/14

Comparative analysis of *S. diclina* Delta 12-desaturase (SDD12.pep)
& *G. hirsutum* Delta 12-desaturase (GH06DES)

Frame: 3 initn: 992 init1: 413 opt: 1086 Z-score: 1531.8 expect(): 2.3e-77
45.6% identity in 379 aa overlap
(9-380:14-384)

SDD12.pep	MCKGQAPSKADVFAAGYRPVAGTPEP-----LPLEPPTITLKDLRAAIPAHC
GH06DESAT	LRVSSTWRXTAFFKASKMGAGGRMPIDGIKEENRGSVNRVPIEKPPFTLGQIKQAIPPHC
	10 20 30 40
SDD12.pep	FERSAATSFYHLAKNLAICAGVFAVGLKLAADLPLAAKLVAWPYWFVQGTFTGIWVI
GH06DESAT	FRRSLLRSFSYVVHDLCLASFFYYIATSYFHF-LPQPFYSYIAPVYVWLQGCILTGVWVI
	50 60 70 80 90 100
SDD12.pep	AHECGHQAFSASEILNDTVGIIHLSLLFVPYHSWKITHRRHHSNTGSCENDEVFTPTPRS
GH06DESAT	AHEWGHHAFRDYQWVDDTVGLILHSALLVPYFSWKISHRRHHSNTGSMERDEVFVPPKPS
	110 120 130 140 150 160
SDD12.pep	VVEAKHDHSLLEESPLYNLYGIVMMLLVGWMPGYLFFNATGPTKYAGLAKSHFNPYAAFF
GH06DESAT	KLSC---FAKYLNNPPGRVLSLVVTLTLGW-PMYLAFFNVSG--RYYDRLASHYNPYGPIY
	170 180 190 200 210 220
SDD12.pep	LPKERLSIWWSLCLFLAALYGFYGVSVFGLLDVARHYIVPYLICNAYLVLITYLQHTDT
GH06DESAT	SDRERLQVYISDTGIFAVIYVLYKIAATKGLAWLLCTYGVPLLIVNAFLVLITYLQHTHS
	230 240 250 260 270 280
SDD12.pep	YVPHFRGDEWNWLRGALCTVDRSFGAWIDSAIHADIADTHVTHHIFSKTPFYHAIEATDAI
GH06DESAT	ALPHYDSSEWDWLRGALSTMDRDFGV-LNKVFHNITDTHVAHHLFSTMPHYHAMEATKAI
	290 300 310 320 330 340
SDD12.pep	TPLLGKYLLIDPTPIPLALWRSFTHCKYVEDDGNVVFYKRKLEEK
GH06DESAT	KPILGKYYPFDGTPIYKAMWREAKECLYVEPDVGGGGGSGKGVFWYRNKFXRPTNCLIAG
	350 360 370 380 390
	1060 1090 1120 1150 1180 1210

FIG.8

9/14

Sequence ID:

Sequence ID 1

5'-ATC CGC GCC GCC ATC CCC AAG CAC TGC TGG GTC AAG-3'

Sequence ID 2

5'-GCC CTC TTC GTC CTC GGC CAY GAC TGC GGC CAY GGC TCG TTC TCG-3'

Sequence ID 3

5'-GAG RTG GTA RTG GGG GAT CTG GGG GAA GAR RTG RTG GRY GAC RTG-3'

Sequence ID 4

5'-CCC TAC CAY GGC TGG CGC ATC TCG CAY CGC ACC CAY CAY CAG AAC-3'

Sequence ID 5

5'-GTT CTG RTG RTG GGT CCG RTG CGA GAT GCG CCA GCC RTG GTA GGG-3'

Sequence ID 6

5'-GGC TCG CAC TTC SAC CCC KAC TCG GAC CTC TTC GTC-3'

Sequence ID 7

5'-GAC GAA GAG GTC CGA GTM GGG GTW GAA GTG CGA GCC-3'

Sequence ID 8

5'-GCG CTG GAK GGT GGT GAG GCC GCC GCG GAW GSA CGA CCA-3'

Sequence ID 9

5'-CTG GGG GAA GAG RTG RTG GAT GAC RTG GGT GCC GAT GTC RTG RTG-3'

Sequence ID 10

5'-GGT GGC CTC GAY GAG RTG GTA RTG GGG GAT CTK GGG GAA GAR RTG-3'

Sequence ID 11

5'-GAG RTG GTA RTG GGG GAT CTG GGG GAA GAR RTG RTG GRY GAC RTG-3'

Sequence ID 12

5'-TAC GCG TAC CTC ACG TAC TCG CTC G-3'

Sequence ID 13

5'-TTC TTG CAC CAC AAC GAC GAA GCG ACG-3'

Sequence ID 14

5'-GGA GTG GAC GTA CGT CAA GGG CAA C-3'

Sequence ID 15

5'-TCA AGG GCA ACC TCT CGA GCG TCG AC-3'

Sequence ID 16

5'-CCC AGT CAC GAC GTT GTA AAA CGA CGG CCA G-3'

FIG.9A

10/14

Sequence ID 17

5'- AGC GGA TAA CAA TTT CAC ACA GGA AAC AGC -3'

Sequence ID 18

5'-GGT AAA AGA TCT CGT CCT TGT CGA TGT TGC-3'

Sequence ID 19

5'-GTC AAA GTG GCT CAT CGT GC-3'

Sequence ID 20

5'-CGA GCG AGT ACG TGA GGT ACG CGT AC-3'

Sequence ID 21

5'-TCA ACA GAA TTC ATG ACC GAG GAT AAG ACG AAG GTC GAG TTC CCG-3'

Sequence ID 22

5'-AAA AGA AAG CTT CGC TTC CTA GTC TTA GTC CGA CTT GGC CTT GGC-3'

Sequence ID 23

5'-TCA ACA AAG CTT ATG ACC GAG GAT AAG ACG AAG GTC GAG TTC CCG-3'

Sequence ID 24

5'-AAA AGA GAA TTC CGC TTC CTA GTC TTA GTC CGA CTT GGC CTT GGC-3'

Sequence ID 25

```

1  ATGACTGAGG ATAAGACGAA GGTGAGTTC CCGACGCTCA CGGAGCTCAA
51  GCACTCGATC CCGAACGCGT GCTTTGAGTC GAACCTCGGC CTCTCGTCT
101 ACTACACGGC CCGCGCGATC TTCAACGCGT CGGCCTCGGC GGCCTGCTC
151 TACGCGGCGC GCTCGACGCC GTTCATTGCC GATAACGTTT TGCTCCACGC
201 GCTCGTTTGC GCCACCTACA TCTACGTGCA GGGCGTCATC TTCTGGGGCT
251 TCTTCACGGT CGGCCACGAC TGGCGGCACT CGGCCTTCTC GCGCTACCAC
301 AGCGTCAACT TTATCATCGG CTGCATCATG CACTCTGCGA TTTTGACGCC
351 GTTCGAGAGC TGGCGCGTGA CGCACCGCCA CCACCACAAG AACACGGGCA
401 ACATTGATAA GGACGAGATC TTTTACCCGC ACCGGTCGGT CAAGGACCTC
451 CAGGACGTGC GCCAATGGGT CTACACGCTC GGCAGTGGT GGTTCGTCTA
501 CTTGAAGGTC GGGTATGCCC CGCGCACGAT GAGCCACTTT GACCCGTGGG
551 ACCCGCTCCT CTTTCGCGGC GCGTCGGCCG TCATCGTGTC GCTCGGCGTC
601 TGGGCGCCTT TCTTCGCGGC GTACGCGTAC CTCACATACT CGCTCGGCTT
651 TGCCGTCATG GGCCTCTACT ACTATGCGCC GCTCTTTGTC TTTGCTTCGT
701 TCCTCGTCAT TACGACCTT TTGCACCACA ACGACGAAGC GACGCCGTGG
751 TACGGCGACT CGGAGTGGAC GTACGTCAAG GGCAACCTCT CGAGCGTCGA
801 CCGCTCGTAC GGCAGGTTTC TGGACAACCT GAGCCACCAC ATTGGCACGC
851 ACCAGGTCCA CCACTTGTTT CCGATCATTG CGCACTACAA GCTCAACGAA
901 GCCACCAAGC ACTTTGCGGC CGCGTACCCG CACCTCGTGC GCAGGAACGA
951 CGAGCCCATC ATCAGGCCTT TCTTCAAGAC CGCGCACCTC TTTGTCAACT
1001 ACGGCGCTGT GCCCGAGACG GCGCAGATCT TCACGCTCAA AGAGTCGGCC
1051 GCGGCGGCCA AGGCCAAGTC GGAATAA

```

FIG.9B

11/14

Sequence ID 26

1 MTEDKTKVEF PTLTEKHSI PNACFESNLG LSLYYTARAI FNASASAALL
51 YAARSTPFIA DNVLLHALVC ATYIYVQGI FWGFFTVGHD CGHSAFSRYH
101 SVNFIIGCIM HSAILTPFES WRVTHRHKK NTGNIDKDEI FYPHRSVKDL
151 QDVRQWVYTL GGAWFVYLKV GYAPRTMSHF DPWDPLLLRR ASAVIVSLGV
201 WAAFFAAYAY LTYSLGFAVM GLYYYAPLFV FASFLVITTF LHNDEATPW
251 YGDSEWTVYK GNLSSVDRSY GAFVDNLSSH IGTHQVHHLF PIIPHYKLINE
301 ATKHFAAYP HLVRNDEPI ITAFFKTAHL FVNYGAVPET AQIFTLKESA
351 AAKAKSD*

Sequence ID 27

1 ATGGCCCCGC AGACGGAGCT CCGCCAGCGC CACGCCGCCG TCGCCGAGAC
51 GCCGGTGGCC GGCAAGAAGG CCTTTACATG GCAGGAGGTC GCGCAGCACA
101 ACACGGCGGC CTCGGCCTGG ATCATTATCC GCGGCAAGGT CTACGACGTG
151 ACCGAGTGGG CCAACAAGCA CCCCAGCGGC CCGGAGATGG TGCTGCTGCA
201 CGCCGGTCGC GAGGCCACCG ACACGTTTGA CTCGTACCAC CCGTTCAGCG
251 ACAAGGCCGA GTCGATCTTG AACAAGTATG AGATTGGCAC GTTCACGGGC
301 CCGTCCGAGT TTCCGACCTT CAAGCCGGAC ACGGGCTTCT ACAAGGAGTG
351 CCGCAAGCGC GTTGGCGAGT ACTTCAAGAA GAACAACCTC CATCCGCGAGG
401 ACGGCTTCCC GGGCCTCTGG CGCATGATGG TCGTGTTCG GGTCCGCGGC
451 CTCGCCCTGT ACGGCATGCA CTTTTGACT ATCTTTGCGC TGCAGCTCGC
501 GGCCGCGGCG CTCTTTGGCG TCTGCCAGGC GCTGCCGCTG CTCACGTCA
551 TGCACGACTC GTCGCACGCG TCGTACACCA ACATGCCGTT CTTCCATTAC
601 GTCGTCCGCC GCTTTGCCAT GGACTGGTTT GCCGCGGCT CGATGGTGTC
651 ATGGCTCAAC CAGCACGTGC TGGGCCACCA CATCTACAG AACGTCCGCG
701 GCTCGGACCC GGATCTTCCG GTCAACATGG ACGGCGACAT CCGCCGATC
751 GTGAACCGCC AGGTGTTCCA GCCCATGTAC GCATTCCAGC ACATCTACCT
801 TCCGCGGCTC TATGGCGTGC TTGGCCTCAA GTTCCGATC CAGGACTTCA
851 CCGACACGTT CGGCTCGCAC ACGAACGGCC CGATCCGCGT CAACCCGCAC
901 GCGCTCTCGA CGTGGATGGC CATGATCAGC TCCAAGTCGT TCTGGGCCCTT
951 CTACCGCGTG TACCTTCCGC TTGCCGTGCT CCAGATGCC ATCAAGACGT
1001 ACCTTGCGAT CTTCTTCTC GCCGAGTTTG TCACGGGCTG GTACCTCGCG
1051 TTCAACTTCC AAGTAAGCCA TGTCTCGACC GAGTGGGCT ACCCATGCGG
1101 CGACGAGGCC AAGATGGCGC TCCAGGACGA GTGGGAGTC TCGCAGGTCA
1151 AGACGTCCGT CCACTACGCC CATGGCTCGT GGATGACGAC GTTCCTTGCC
1201 GGCGCGCTCA ACTACCAGGT CGTGCACCAC TTGTTCCCA GCGTGTGCGA
1251 GTACCACTAC CCGGCGATCG CGCCCATCAT CGTCGACGTC TGCAAGGAGT
1301 ACAACATCAA GTACGCCATC TTGCCGACT TTACGGCGGC GTTCGTTGCC
1351 CACTTGAAGC ACCTCCGCAA CATGGGCCAG CAGGGCATCG CCGCCACGAT
1401 CCACATGGGC TAA

FIG.9C

12/14

Sequence ID 28

```

1  ATGGCAAACA GCAGCGTGTG GGATGATGTG GTGGGCCGCG TGGAGACCGG
51  CGTGGACCAG TGGATGGATG GCGCCAAGCC GTACGCACTC ACCGATGGGC
101 TCCCGATGAT GGACGTGTCC ACCATGCTGG CATTGAGGT GGGATACATG
151 GCCATGCTGC TCTTCGGCAT CCCGATCATG AAGCAGATGG AGAAGCCTTT
201 TGAGCTCAAG ACCATCAAGC TCTTGACAA CTTGTTTCTC TTCGGACTTT
251 CCTTGTACAT GTGCGTGGAG ACCATCCGCC AGGCTATCCT CGGAGGCTAC
301 AAAGTGTTTG GAAACGACAT GGAGAAGGGC AACGAGTCTC ATGCTCAGGG
351 CATGTCTCGC ATCGTGTACG TGTCTGCGT GTCCAAGGCA TACGAGTTCT
401 TGGATACCGC CATCATGATC CTTTGCAAGA AGTTCAACCA GGTTTCCTTC
451 TTGCATGTGT ACCACCATGC CACCATTTTT GCCATCTGGT GGGCTATCGC
501 CAAGTACGCT CCAGGAGGTG ATGCGTACTT TTCAGTGATC CTCAACTCTT
551 TCGTGACAC CGTCATGTAC GCATACTACT TCTTCTCCTC CCAAGGGTTC
601 GGGTTCGTGA AGCCAATCAA GCCGTACATC ACCACCCTTC AGATGACCCA
651 GTTCATGGCA ATGCTTGTGC AGTCCTTGTA CGACTACCTC TTCCCATGCG
701 ACTACCCACA GGCTCTTGTG CAGCTTCTTG GAGTGTACAT GATCACCTTG
751 CTTGCCCTCT TCGGCAACTT TTTTGTGCAG AGCTATCTTA AAAAGCCAAA
801 AAAGAGCAAG ACCAACTAA

```

Sequence ID 29

```

1  MTVGFDETVT MDTVRNHNMP DDAWCAIHGT VYDITKFSKV HPGGDIIMLA
51  AGKEATILFE TYHIKGVPA VLRKYKVGKL PQGKKGETSH MPTGLDSASY
101 YSWDSEFYRV LRERVAKKLA EPGLMQRARM ELWAKAIFLL AGFWGSLYAM
151 CVLDPHGGAM VAAVTLGVFA AFVGTCTIQHD GSHGAFSKSR FMNKAAGWTL
201 DMIGASAMTW EMQHVLGHP YTNLIEMENG LAKVKGADVD PKKVDQESDP
251 DVFSTYMLR LHPWHRQRFY HKFQHLYAPL IFGFMINKV ISQDVGVVLR
301 KRLFQIDANC RYGSPWNVAR FWIMKLLTTL YMVALPMYMQ GPAQGLKLFF
351 MAHFTCGEVL ATMFIVNHII EGVSYASKDA VKGVMAPPRT VHGVTPMQVT
401 QKALSAAEST KSDADKTTMI PLNDWAAVQC QTSVNWAVGS WFWNHFSGGL
451 NHQIEHHCFP QNPHTVNVYI SGIVKETCEE YGVPIQAEIS LFSAYFKMLS
501 HLRTLGNEDL TAWST*

```

Sequence ID 30

5'-CCG SAG TTC ACS ATC AAG GAG ATC CGC GAS KSC ATC CCG GCC CAC TGC TTC -3'

Sequence ID 31

5'-GRS CTT CTT GAK GTG GWM SGT GGC CTC CTC GGC GTG GTA GWR CGG CAT-3'

Sequence ID 32

5'-CCS STC TAC TGG GCC TGC CAG GGT RTC GTC CTC ACS GGT GTC TGG-3'

FIG.9D

13/14

Sequence ID 33

5'-CCS STC TAC TGG ATC RYS CAG GGT RTC GTC KGY ACS GGT GTC TGG-3'

Sequence ID 34

5'-GGC GTG GTA GTG CGG CAT SMM CGA GAA GAR GTG GTG GGC GAC GTG-3'

Sequence ID 35

5'-CAC GTA CCT CCA GCA CAC GGA CAC CTA CG-3'

Sequence ID 36

5'- GAT CGA CAG CGC GAT CCA CCA CAT TGC-3'

Sequence ID 37

5'- CAA ATG GTA AAA GCT AGT GGC AGC GCT GC-3'

Sequence ID 38

5'-AGT ACG TGC CCT GGA CGA ACC AGT AGA TG-3'

Sequence ID 39

5'- TCA ACA GAA TTC ATG TGC AAA GGT CAA GCT CCT TCC AAG GCC GAC GTG -3'

Sequence ID 40

5'- AAA AGA AAG CTT TTA CTT TTC CTC GAG CTT GCG CTT GTA AAA CAC AAC-3'

Sequence ID 41

```

1  ATGTGCAAAG GTCAAGCTCC TTCCAAGGCC GACGTGTTCC ACGCTGCGGG
51  GTACCGCCCG GTCGCCGGCA CGCCCGAGCC GCTGCCGCTG GAGCCCCCGA
101 CGATCACGCT CAAGGACCTG CGCGCGGCGA TCCCGGCCCA CTGCTTTGAG
151 CGCAGCGCTG CCACTAGCTT TTACCATTG GCCAAGAACC TTGCGATCTG
201 CGCCGGCGTG TTCGCCGTTG GCCTCAAGCT CGCGGCTGCC GACTTGCCGC
251 TCGCGGCCAA GCTGGTCGCG TGGCCCATCT ACTGGTTCGT CCAGGGCAGC
301 TACTTTACGG GCATCTGGGT CATTGCGCAC GAATGCGGCC ACCAGGCGTT
351 CTCGGCGTCC GAGATCCTCA ACGACACGGT CGGTATCATT CTTCACTCGC
401 TCCTCTTTGT GCCGTACCAC AGCTGGAAGA TCACGCACCG CCGCCACCAC
451 TCCAACACGG GCAGCTGCGA GAACGACGAG GTGTTTACGC CGACGCCGCG
501 GTCCGTCGTC GAGGCCAAGC ACGACCACTC GTCCTCGAA GAGAGCCCGC
551 TCTACAACT GTACGGCATC GTCATGATGC TTCTCGTGGG CTGGATGCCG
601 GGCTACCTCT TCTTCAACGC GACCGGCCCG ACCAAGTACG CTGGCCTCGC
651 CAAGTCGCAC TTCAACCCGT ACGCAGCCTT TTTCTCCCA AAGGAGCGCC
701 TCAGCATCTG GTGGAGCGAC CTCTGCTTCC TCGCGGCTT GTACGGCTTT
751 GGCTACGCG TCTCGGTCTT CGGCCTCCTC GATGTCGCCC GCCACTACAT
801 CGTGCCGTAC CTCATTTGCA ACGCGTACCT CGTGCTCATC ACGTACCTCC
851 AGCACACGGA TACGTACGTG CCCCATTCC GCGGCGACGA GTGGAACCTGG
901 CTGCGCGGCG CGCTCTGCAC CGTCGACCGC TCGTTCGGCG CGTGGATCGA
951 CAGCGCGATC CACCACATTG CCGACACGCA CGTGACGCAC CACATTTTCT
1001 CCAAGACGCC CTTCTACCAC GCGATCGAGG CGACCGACGC CATCACGCCC
1051 CTCCTCGGCA AGTACTACCT CATCGACCGG ACGCCGATCC CGCTGGCGCT
1101 CTGGCGCTCG TTCACGCACT GCAAGTACGT CGAGGACGAC GGCAACGTTG
1151 TGTTTTACAA GCGCAAGCTC GAGGAAAAGT AA

```

FIG.9E

14/14

Sequence ID 42

1 MCKGQAPSKA DVFHAAGYRP VAGTPEPLPL EPPTITLKDL RAAIPAHCFE
51 RSAATSFYHL AKNLAICAGV FAVGLKLAAL DLPLAAKLVA WPIYWFVQGT
101 YFTGIWVIAH ECGHQAFSAS EILNDTVGII LHSLLFVPYH SWKITHRRHH
151 SNTGSCENDE VFTPTPRSVV EAKHDHSLLE ESPLYNLYGI VMMLLVGWMP
201 GYLFFNATGP TKYAGLAKSH FNPYAAFFLP KERLSIWWSL LCFLAALYGF
251 GYGVSFVGLL DVARHYIVPY LICNAYLVLI TYLQHTDTYV PHFRGDEWNW
301 LRGALECTVDR SFGAWIDSAI HHIADTHVTH HIFSKTPFYH AIEATDAITP
351 LLGKYYLIDP TPIPLALWRS FTHCKYVEDD GNVVFYKRKL EEK*

FIG.9F

SEQUENCE LISTING

<110> Abbott Laboratories
Mukerji, Pradip
Huang, Yung-Sheng
Pereira, Suzette L.

<120> DESATURASE GENES, ENZYMES ENCODED
THEREBY, AND USES THEREOF

<130> 6884.US.O1

<140> 10/060,793

<141> 2002-01-30

<160> 60

<170> FastSEQ for Windows Version 4.0

<210> 1

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Forward Primer R01144

<400> 1

atccgcgcgcg ccatccccaa gcactgctgg gtcaag

36

<210> 2

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Forward Primer R01119

<221> misc_feature

<222> (21)...(21)

<223> y = t/u or c at position 21

<221> misc_feature

<222> (33)...(33)

<223> y = t/u or c at position 33

<400> 2

gccctcttcg tcctcggcca ygactgcggc cayggctcgt tctcg

45

<210> 3

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Reverse Primer R01118

<221> misc_feature
<222> (4)...(4)
<223> r = g or a at position 4

<221> misc_feature
<222> (10)...(10)
<223> r = g or a at position 10

<221> misc_feature
<222> (30)...(31)
<223> r = g or a at positions 30-31

<221> misc_feature
<222> (34)...(34)
<223> r = g or a at position 34

<221> misc_feature
<222> (38)...(38)
<223> r = g or a at position 38

<221> misc_feature
<222> (39)...(39)
<223> y = t/u or c at position 39

<221> misc_feature
<222> (43)...(43)
<223> r = g or a at position 43

<400> 3
gagrtggtar tgggggatct gggggaagar rtgrtggrtg acrtg

<210> 4
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Forward Primer RO1121

<221> misc_feature
<222> (9)...(9)
<223> y = t/u or c at position 9

<221> misc_feature
<222> (27)...(27)
<223> y = t/u or c at position 27

<221> misc_feature
<222> (36)...(36)
<223> y = t/u or c at position 36

<221> misc_feature
<222> (39)...(39)
<223> y = t/u or c at position 39

<400> 4

ccctaccayg gctggcgcat ctcgcaycgc acccaycayc agaac

45

<210> 5
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Reverse Primer R01122

<221> misc_feature
<222> (7)...(7)
<223> r = g or a at position 7

<221> misc_feature
<222> (10)...(10)
<223> r = g or a at position 10

<221> misc_feature
<222> (37)...(37)
<223> r = g or a at position 37

<400> 5
gttctgrtgr tgggtccgrt gcgagatgcg ccagccrtgg taggg

45

<210> 6
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Forward Primer R01146

<221> misc_feature
<222> (13)...(13)
<223> s = g or c at position 13

<221> misc_feature
<222> (19)...(19)
<223> k = g or t/u at position 19

<400> 6
ggctcgact tcsaccccka ctcggacctc ttcgtc

36

<210> 7
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Reverse Primer R01147

<221> misc_feature
<222> (18)...(18)
<223> m = a or c at position 18

<221> misc_feature

<222> (24)...(24)
<223> w = a or t/u at position 24

<400> 7
gacgaagagg tccgagtmgg ggtwgaagtg cgagcc

36

<210> 8
<211> 39
<212> DNA
<213> Artificial Sequence

<220>
<223> Reverse Primer R01148

<221> misc_feature
<222> (9)...(9)
<223> k = g or t/u at position 9

<221> misc_feature
<222> (30)...(30)
<223> w = a or t/u at position 30

<221> misc_feature
<222> (32)...(32)
<223> s = g or c at position 32

<400> 8
gcgctggakg gtggtgaggc cgccgcggaw gsacgacca

39

<210> 9
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Reverse Primer R01114

<221> misc_feature
<222> (13)...(13)
<223> r = g or a at position 13

<221> misc_feature
<222> (16)...(16)
<223> r = g or a at position 16

<221> misc_feature
<222> (25)...(25)
<223> r = g or a at position 25

<221> misc_feature
<222> (40)...(40)
<223> r = g or a at position 40

<221> misc_feature
<222> (43)...(43)
<223> r = g or a at position 43

<400> 9
ctgggggaag agrtgrtgga tgacrtgggt gccgatgtcr tgrtg

45

<210> 10
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Reverse Primer R01116

<221> misc_feature
<222> (12)...(12)
<223> y = t/u or c at position 12

<221> misc_feature
<222> (16)...(16)
<223> r = g or a at position 16

<221> misc_feature
<222> (22)...(22)
<223> r = g or a at position 22

<221> misc_feature
<222> (33)...(33)
<223> k = g or t/u at position 33

<221> misc_feature
<222> (42)...(43)
<223> r = g or at at positions 42-43

<400> 10
ggtggcctcg aygagrtggt artgggggat ctkggggaag arrtg

45

<210> 11
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Reverse Primer R01118

<221> misc_feature
<222> (4)...(4)
<223> r = g or a at position 4

<221> misc_feature
<222> (10)...(10)
<223> r = g or a at position 10

<221> misc_feature
<222> (30)...(31)
<223> r = g or a at positions 30-31

<221> misc_feature
<222> (34)...(34)
<223> r = g or a at position 34

6.

<221> misc_feature
<222> (38)...(38)
<223> r = g or a at position 38

<221> misc_feature
<222> (39)...(39)
<223> y = t/u or c at position 39

<221> misc_feature
<222> (43)...(43)
<223> r = g or a at position 43

<400> 11
gagrtggtar tgggggatct gggggaagar rtgrtggryg acrtg

45

<210> 12
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Forward Primer RO1188

<400> 12
tacgcgtacc tcacgtactc gctcg

25

<210> 13
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Forward Primer RO1189

<400> 13
ttcttgcacc acaacgacga agcgacg

27

<210> 14
<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> Forward Primer RO1190

<400> 14
ggagtggacg tacgtcaagg gcaac

25

<210> 15
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Forward Primer RO1191

<400> 15
tcaagggcaa cctctcgagc gtcgac 26

<210> 16
<211> 31
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer RO898

<400> 16
cccagtcacg acgttgtaaa acgacggcca g 31

<210> 17
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Primer RO899

<400> 17
agcggataac aatttcacac aggaaacagc 30

<210> 18
<211> 30
<212> DNA
<213> Artificial Sequence

<220>
<223> Reverse Primer RO1185

<400> 18
ggtaaaagat ctcgtccttg tcgatgttgc 30

<210> 19
<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Reverse Primer RO1186

<400> 19
gtcaaagtgg ctcacgtgc 20

<210> 20
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> Reverse Primer RO1187

<400> 20
cgagcgagta cgtgaggtac gcgtac 26

<210> 21
 <211> 45
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Forward Primer RO1212

<400> 21
 tcaacagaat tcatgaccga ggataagacg aaggtcgagt tcccg 45

<210> 22
 <211> 45
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Reverse Primer RO1213

<400> 22
 aaaagaaagc ttcgcttcct agtcttagtc cgacttggcc ttggc 45

<210> 23
 <211> 45
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Forward Primer RO1221

<400> 23
 tcaacaaagc ttatgaccga ggataagacg aaggtcgagt tcccg 45

<210> 24
 <211> 45
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Reverse Primer RO1222

<400> 24
 aaaagagaat tccgcttcct agtcttagtc cgacttggcc ttggc 45

<210> 25
 <211> 1077
 <212> DNA
 <213> Saprolegnia diclina

<400> 25
 atgactgagg ataagacgaa ggtcgagttc ccgacgctca cggagctcaa gcactcgatc 60
 ccgaacgcgt gctttgagtc gaacctcggc ctctcgctct actacacggc ccgcgcgatc 120
 ttcaacgcgt cggcctcggc ggcgtgctc tacgcggcgc gctcgacgcc gttcattgcc 180
 gataacgttc tgctccacgc gctcgtttgc gccacctaca tctacgtgca gggcgatc 240
 ttctggggct tcttcacggt cggccacgac tgccggccact cggccttctc gcgctaccac 300
 agcgtcaact ttatcatcgg ctgcatcatg cactctgcga ttttgacgcc gttcgagagc 360

```

tggcgcgtga cgcaccgcca ccaccacaag aacacgggca acattgataa ggacgagatc 420
ttttaccgcg accggtcggt caaggacctc caggacgtgc gccaatgggt ctacacgctc 480
ggcgggtgcgt ggtttgtcta cttgaaggtc gggatatgcc cgcgcacgat gagccacttt 540
gacccggtggg acccgctcct ccttcgccgc gcgtcggccg tcatcgtgtc gctcggcgtc 600
tgggccgcct tcttcgccgc gtacgcgtac ctcacatact cgctcggctt tgccgtcatg 660
ggcctctact actatgcgcc gctctttgtc tttgcttcgt tcctcgtcat tacgaccttc 720
ttgcaccaca acgacgaagc gacgccgtgg tacggcgact cggagtggac gtacgtcaag 780
ggcaacctct cgagcgtcga ccgctcgtac ggcgcggttcg tggacaacct gagccaccac 840
attggcacgc accaggtcca ccacttggtc ccgatcattc cgactacaa gctcaacgaa 900
gccaccaagc actttgcggc cgcgtaaccg cacctcgtgc gcaggaacga cgagcccatc 960
atcacggcct tcttcaagac cgcgcacctc tttgtcaact acggcgctgt gcccgagacg 1020
gcgcagatct tcacgctcaa agagtcggcc gcggccgcca aggccaagtc ggactaa 1077

```

<210> 26

<211> 358

<212> PRT

<213> Saprolegnia diclina

<400> 26

```

Met Thr Glu Asp Lys Thr Lys Val Glu Phe Pro Thr Leu Thr Glu Leu
1      5      10      15
Lys His Ser Ile Pro Asn Ala Cys Phe Glu Ser Asn Leu Gly Leu Ser
20      25      30
Leu Tyr Tyr Thr Ala Arg Ala Ile Phe Asn Ala Ser Ala Ser Ala Ala
35      40      45
Leu Leu Tyr Ala Ala Arg Ser Thr Pro Phe Ile Ala Asp Asn Val Leu
50      55      60
Leu His Ala Leu Val Cys Ala Thr Tyr Ile Tyr Val Gln Gly Val Ile
65      70      75      80
Phe Trp Gly Phe Phe Thr Val Gly His Asp Cys Gly His Ser Ala Phe
85      90      95
Ser Arg Tyr His Ser Val Asn Phe Ile Ile Gly Cys Ile Met His Ser
100      105      110
Ala Ile Leu Thr Pro Phe Glu Ser Trp Arg Val Thr His Arg His His
115      120      125
His Lys Asn Thr Gly Asn Ile Asp Lys Asp Glu Ile Phe Tyr Pro His
130      135      140
Arg Ser Val Lys Asp Leu Gln Asp Val Arg Gln Trp Val Tyr Thr Leu
145      150      155      160
Gly Gly Ala Trp Phe Val Tyr Leu Lys Val Gly Tyr Ala Pro Arg Thr
165      170      175
Met Ser His Phe Asp Pro Trp Asp Pro Leu Leu Leu Arg Arg Ala Ser
180      185      190
Ala Val Ile Val Ser Leu Gly Val Trp Ala Ala Phe Phe Ala Ala Tyr
195      200      205
Ala Tyr Leu Thr Tyr Ser Leu Gly Phe Ala Val Met Gly Leu Tyr Tyr
210      215      220
Tyr Ala Pro Leu Phe Val Phe Ala Ser Phe Leu Val Ile Thr Thr Phe
225      230      235
Leu His His Asn Asp Glu Ala Thr Pro Trp Tyr Gly Asp Ser Glu Trp
245      250      255
Thr Tyr Val Lys Gly Asn Leu Ser Ser Val Asp Arg Ser Tyr Gly Ala
260      265      270
Phe Val Asp Asn Leu Ser His His Ile Gly Thr His Gln Val His His
275      280      285
Leu Phe Pro Ile Ile Pro His Tyr Lys Leu Asn Glu Ala Thr Lys His
290      295      300

```

Phe Ala Ala Ala Tyr Pro His Leu Val Arg Arg Asn Asp Glu Pro Ile
 305 310 315 320
 Ile Thr Ala Phe Phe Lys Thr Ala His Leu Phe Val Asn Tyr Gly Ala
 325 330 335
 Val Pro Glu Thr Ala Gln Ile Phe Thr Leu Lys Glu Ser Ala Ala Ala
 340 345 350
 Ala Lys Ala Lys Ser Asp
 355

<210> 27
 <211> 1413
 <212> DNA
 <213> *Saprolegnia diclina*

<400> 27
 atggccccgc agacggagct ccgccagcgc cagccgcgcg tcgccgagac gccggtggcc 60
 ggcaagaagg cctttacatg gcaggaggtc gcgcagcaca acacggcggc ctcggcctgg 120
 atcattatcc gcggcaaggt ctacgacgtg accgagtggg ccaacaagca cccggcgggc 180
 cgcgagatgg tgctgctgca cgcgggtcgc gaggccaccg acacgttcga ctcgtaccac 240
 ccgttcagcg acaaggccga gtcgatcttg aacaagtatg agattggcac gttcacgggc 300
 ccgtccgagt ttccgacctt caagccggac acgggcttct acaaggagtg ccgcaagcgc 360
 gttggcgagt acttcaagaa gaacaacctc catccgcagg acggcttccc gggcctctgg 420
 cgcattgatgg tcgtgtttgc ggtcgcgggc ctgcgcttgt acggcatgca cttttcgact 480
 atcctttgcgc tgcagctcgc ggccgcgggc ctctttggcg tctgccaggc gctgccgctg 540
 ctccacgtca tgcacgactc gtcgcacgcg tcgtacacca acatgccgtt cttccattac 600
 gtcgtcggcc gctttgccat ggactggttt gccggcggct cgatgggtgc atgggtcaac 660
 cagcacgtcg tgggccacca catctacacg aacgtcgcgc gtcgggaccc ggatcttccg 720
 gtcaacatgg acggcgacat ccgccgcgac gtgaaccgcc aggtgttcca gcccatgtac 780
 gcattccagc acatctacct tccgcgcgtc tatggcgtgc ttggcctcaa gttccgcac 840
 caggacttca ccgacacgtt cggctcgcac acgaacggcc cgatccgcgt caaccgcac 900
 gcgctctcga cgtggatggc catgatcagc tccaagtcgt tctgggcctt ctaccgcgtg 960
 taccttccgc ttgccgtgct ccagatgccc atcaagacgt accttgcgat cttcttcctc 1020
 gccgagtttg tcacgggctg gtacctcgcg ttcaacttcc aagtaagcca tgtctcgacc 1080
 gagtgcggct acccatgcgg cgacgaggcc aagatggcgc tccaggacga gtgggcagtc 1140
 tcgcaggtca agacgtcggg cgactacgcc catggctcgt ggatgacgac gttccttgcc 1200
 ggcgcgctca actaccaggt cgtgcaccac ttgttcccca gcgtgtcgca gtaccactac 1260
 ccggcgatcg cgcccatcat cgtcgacgtc tgcaaggagt acaacatcaa gtacgccatc 1320
 ttgccggact ttacggcggc gttcgttgcc cacttgaagc acctccgcaa catgggcccag 1380
 cagggcatcg ccgccacgat ccacatgggc taa 1413

<210> 28
 <211> 819
 <212> DNA
 <213> *Thraustochytrid* sp.

<400> 28
 atggcaaaaca gcagcgtgtg ggatgatgtg gtgggcccgc tggagaccgg cgtggaccag 60
 tggatggatg gcgccaagcc gtacgcactc accgatgggc tcccgatgat ggacgtgtcc 120
 accatgctgg cattcgaggt gggatacatg gccatgctgc tcttcggcat cccgatcatg 180
 aagcagatgg agaagccttt tgagctcaag accatcaagc tcttgcaaaa cttgtttctc 240
 ttcggacttt ccttgtagat gtgcgtggag accatccgcc aggtatcct cggaggctac 300
 aaagtgtttg gaaacgacat ggagaagggc aacgagtctc atgctcaggg catgtctcgc 360
 atcgtgtacg tgttctgcgt gtccaaggca tacgagttct tggataccgc catcatgatc 420
 ctttgcaaga agttcaacca ggtttccttc ttgcatgtgt accaccatgc caccattttt 480
 gccatctggg gggctatcgc caagtacgct ccaggagggt atgcgtactt ttcagtgatc 540
 ctcaactctt tcgtgcacac cgtcatgtac gcatactact tcttctcctc ccaagggttc 600
 gggttcgtga agccaatcaa gccgtacatc accacccttc agatgaccca gttcatggca 660

atgcttgtgc agtccttgta cgactacctc ttcccatgcg actaccacaca ggctcttgtg	720
cagcttcttg gagtgatcat gatcaccttg cttgccctct tcggcaactt ttttgtgcag	780
agctatctta aaaagccaaa aaagagcaag accaactaa	819

<210> 29
 <211> 515
 <212> PRT
 <213> Saprolegnia diclina

<400> 29

Met Thr Val Gly Phe Asp Glu Thr Val Thr Met Asp Thr Val Arg Asn	
1 5 10 15	
His Asn Met Pro Asp Asp Ala Trp Cys Ala Ile His Gly Thr Val Tyr	
20 25 30	
Asp Ile Thr Lys Phe Ser Lys Val His Pro Gly Gly Asp Ile Ile Met	
35 40 45	
Leu Ala Ala Gly Lys Glu Ala Thr Ile Leu Phe Glu Thr Tyr His Ile	
50 55 60	
Lys Gly Val Pro Asp Ala Val Leu Arg Lys Tyr Lys Val Gly Lys Leu	
65 70 75 80	
Pro Gln Gly Lys Lys Gly Glu Thr Ser His Met Pro Thr Gly Leu Asp	
85 90 95	
Ser Ala Ser Tyr Tyr Ser Trp Asp Ser Glu Phe Tyr Arg Val Leu Arg	
100 105 110	
Glu Arg Val Ala Lys Lys Leu Ala Glu Pro Gly Leu Met Gln Arg Ala	
115 120 125	
Arg Met Glu Leu Trp Ala Lys Ala Ile Phe Leu Leu Ala Gly Phe Trp	
130 135 140	
Gly Ser Leu Tyr Ala Met Cys Val Leu Asp Pro His Gly Gly Ala Met	
145 150 155 160	
Val Ala Ala Val Thr Leu Gly Val Phe Ala Ala Phe Val Gly Thr Cys	
165 170 175	
Ile Gln His Asp Gly Ser His Gly Ala Phe Ser Lys Ser Arg Phe Met	
180 185 190	
Asn Lys Ala Ala Gly Trp Thr Leu Asp Met Ile Gly Ala Ser Ala Met	
195 200 205	
Thr Trp Glu Met Gln His Val Leu Gly His His Pro Tyr Thr Asn Leu	
210 215 220	
Ile Glu Met Glu Asn Gly Leu Ala Lys Val Lys Gly Ala Asp Val Asp	
225 230 235 240	
Pro Lys Lys Val Asp Gln Glu Ser Asp Pro Asp Val Phe Ser Thr Tyr	
245 250 255	
Pro Met Leu Arg Leu His Pro Trp His Arg Gln Arg Phe Tyr His Lys	
260 265 270	
Phe Gln His Leu Tyr Ala Pro Leu Ile Phe Gly Phe Met Thr Ile Asn	
275 280 285	
Lys Val Ile Ser Gln Asp Val Gly Val Val Leu Arg Lys Arg Leu Phe	
290 295 300	
Gln Ile Asp Ala Asn Cys Arg Tyr Gly Ser Pro Trp Asn Val Ala Arg	
305 310 315 320	
Phe Trp Ile Met Lys Leu Leu Thr Thr Leu Tyr Met Val Ala Leu Pro	
325 330 335	
Met Tyr Met Gln Gly Pro Ala Gln Gly Leu Lys Leu Phe Phe Met Ala	
340 345 350	
His Phe Thr Cys Gly Glu Val Leu Ala Thr Met Phe Ile Val Asn His	
355 360 365	
Ile Ile Glu Gly Val Ser Tyr Ala Ser Lys Asp Ala Val Lys Gly Val	

12

```

      370      375      380
Met Ala Pro Pro Arg Thr Val His Gly Val Thr Pro Met Gln Val Thr
385      390      395
Gln Lys Ala Leu Ser Ala Ala Glu Ser Thr Lys Ser Asp Ala Asp Lys
      405      410      415
Thr Thr Met Ile Pro Leu Asn Asp Trp Ala Ala Val Gln Cys Gln Thr
      420      425      430
Ser Val Asn Trp Ala Val Gly Ser Trp Phe Trp Asn His Phe Ser Gly
      435      440      445
Gly Leu Asn His Gln Ile Glu His His Cys Phe Pro Gln Asn Pro His
      450      455      460
Thr Val Asn Val Tyr Ile Ser Gly Ile Val Lys Glu Thr Cys Glu Glu
465      470      475
Tyr Gly Val Pro Tyr Gln Ala Glu Ile Ser Leu Phe Ser Ala Tyr Phe
      485      490      495
Lys Met Leu Ser His Leu Arg Thr Leu Gly Asn Glu Asp Leu Thr Ala
      500      505      510
Trp Ser Thr
      515

```

<210> 30

<211> 51

<212> DNA

<213> Artificial Sequence

<220>

<223> Forward Primer RO967

<221> misc_feature

<222> (4)...(4)

<223> s = g or c at position 4

<221> misc_feature

<222> (12)...(12)

<223> s = g or c at position 12

<221> misc_feature

<222> (30)...(30)

<223> s = g or c at position 30

<221> misc_feature

<222> (31)...(31)

<223> k = g or t/u at position 31

<221> misc_feature

<222> (32)...(32)

<223> s = g or c at position 32

<400> 30

ccgsagttca csatcaagga gatccgcgas kscatcccgg cccactgctt c

51

<210> 31

<211> 48

<212> DNA

<213> Artificial Sequence

<220>

13

<223> Reverse Primer R0968

<221> misc_feature

<222> (2)...(2)

<223> r = g or a at position 2

<221> misc_feature

<222> (3)...(3)

<223> s = g or c at position 3

<221> misc_feature

<222> (12)...(12)

<223> k = g or t/u at position 12

<221> misc_feature

<222> (17)...(17)

<223> w = a or t/u at position 17

<221> misc_feature

<222> (18)...(18)

<223> m = a or c at position 18

<221> misc_feature

<222> (19)...(19)

<223> s = g or c at position 19

<221> misc_feature

<222> (41)...(41)

<223> w = a or t/u at position 41

<221> misc_feature

<222> (42)...(42)

<223> r = g or a at position 42

<400> 31

grscttcttg akgtggwmsg tggcctcctc ggcgtggtag wrcggcat

48

<210> 32

<211> 45

<212> DNA

<213> Artificial Sequence

<220>

<223> Forward Primer R0964

<221> misc_feature

<222> (3)...(4)

<223> s = g or c at positions 3-4

<221> misc_feature

<222> (25)...(25)

<223> r = g or a at position 25

<221> misc_feature

<222> (36)...(36)

<223> s = g or c at position 36

<400> 32
ccsstctact gggcctgccca gggtrtcgtc ctcacsggtg tctgg

45

<210> 33
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Forward Primer R0965

<221> misc_feature
<222> (3)...(4)
<223> s = g or c at positions 3-4

<221> misc_feature
<222> (16)...(16)
<223> r = g or a at position 16

<221> misc_feature
<222> (17)...(17)
<223> y = t/u or c at position 17

<221> misc_feature
<222> (18)...(18)
<223> s = g or c at position 18

<221> misc_feature
<222> (25)...(25)
<223> r = g or a at position 25

<221> misc_feature
<222> (31)...(31)
<223> k = g or t/u at position 31

<221> misc_feature
<222> (33)...(33)
<223> y = t/u or c at position 33

<221> misc_feature
<222> (36)...(36)
<223> s = g or c at position 36

<400> 33
ccsstctact ggatcrysca gggtrtcgtc kgyacsggtg tctgg

45

<210> 34
<211> 45
<212> DNA
<213> Artificial Sequence

<220>
<223> Reverse Primer R0966

<221> misc_feature
<222> (19)...(19)
<223> s = g or c at position 19

<221> misc_feature
<222> (20)...(21)
<223> m = a or c at positions 20-21

<221> misc_feature
<222> (30)...(30)
<223> r = g or a at position 30

<400> 34
ggcgtggttag tgcggcatsm mcgagaagar gtggtgggacg acgtg 45

<210> 35
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Forward Primer R0975

<400> 35
cacgtacctc cagcacacgg acacctacg 29

<210> 36
<211> 27
<212> DNA
<213> Artificial Sequence

<220>
<223> Forward Primer R0976

<400> 36
gatcgacagc gcgatccacc acattgc 27

<210> 37
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Reverse Primer R0977

<400> 37
caaatggtaa aagctagtgg cagcgctgc 29

<210> 38
<211> 29
<212> DNA
<213> Artificial Sequence

<220>
<223> Reverse Primer R0978

<400> 38
agtacgtgcc ctggacgaac cagtagatg 29

<210> 39

16

<211> 48
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Forward Primer RO1051

<400> 39
 tcaacagaat tcatgtgcaa aggtcaagct ccttccaagg ccgacgtg 48

<210> 40
 <211> 48
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Reverse Primer RO1057

<400> 40
 aaaagaaagc ttttactttt cctcgagctt gcgcttgtaa aacacaac 48

<210> 41
 <211> 1182
 <212> DNA
 <213> Saprolegnia diclina

<400> 41
 atgtgcaaaag gtcaagctcc ttccaaggcc gacgtgttcc acgctgcggg gtaccgcccg 60
 gtcgccggca cgcccgagcc gctgccgctg gagccccga cgatcacgct caaggacctg 120
 cgcgcggcga tcccggccca ctgctttgag cgcagcgctg ccaactagctt ttaccatttg 180
 gccaagaacc ttgcgatctg cgcggcgctg ttccgcttg gcctcaagct cgcggctgcc 240
 gacttgccgc tcgcggccaa gctggtcgctg tggcccatct actggttcgt ccagggcacg 300
 tactttacgg gcatctgggt cattgcgcac gaatgcggcc accaggcggt ctcggcgtcc 360
 gagatcctca acgacacggt cggatatcatt cttcactcgc tcctctttgt gccgtaccac 420
 agctggaaga tcacgcaccg ccgccaccac tccaacacgg gcagctgcga gaacgacgag 480
 gtgtttacgc cgacgcgcgc gtcgcgtcgtc gaggccaagc acgaccactc gtcctcga 540
 gagagcccgc tctacaacct gtacggcatc gtcatgatgc ttctcgtggg ctggatgccg 600
 ggctacctct tcttcaacgc gaccggccc accaagtacg ctggcctcgc caagtcgcac 660
 ttcaaccctt acgcagcctt tttcctccca aaggagcgcc tcagcatctg gtggagcgac 720
 ctctgcttcc tcgcggcctt gtacggcttt ggctacggcg tctcgggtctt cggcctctc 780
 gatgtcgcgc gccactacat cgtgccgtac ctcatattgca acgcgtacct cgtgctcatc 840
 acgtacctcc agcacacgga tacgtacgtg cccacttcc gcggcgacga gtggaactgg 900
 ctgcgcggcg cgctctgcac cgtcgaccgc tcgttcggcg cgtggatcga cagcgcgatc 960
 caccacattg ccgacacgca cgtgacgcac cacattttct ccaagacgcc cttctaccac 1020
 gcgatcgagg cgaccgacgc catcacgccc ctccctcgga agtactacct catcgaccgc 1080
 acgccgatcc cgctggcgct ctggcgctcg ttcacgcact gcaagtacgt cgaggacgac 1140
 ggcaacgctg tgttttataa gcgcaagctc gaggaaaagt aa 1182

<210> 42
 <211> 393
 <212> PRT
 <213> Saprolegnia diclina

<400> 42
 Met Cys Lys Gly Gln Ala Pro Ser Lys Ala Asp Val Phe His Ala Ala
 1 5 10 15
 Gly Tyr Arg Pro Val Ala Gly Thr Pro Glu Pro Leu Pro Leu Glu Pro

20 25 30
 Pro Thr Ile Thr Leu Lys Asp Leu Arg Ala Ala Ile Pro Ala His Cys
 35 40 45
 Phe Glu Arg Ser Ala Ala Thr Ser Phe Tyr His Leu Ala Lys Asn Leu
 50 55 60
 Ala Ile Cys Ala Gly Val Phe Ala Val Gly Leu Lys Leu Ala Ala Ala
 65 70 75 80
 Asp Leu Pro Leu Ala Ala Lys Leu Val Ala Trp Pro Ile Tyr Trp Phe
 85 90 95
 Val Gln Gly Thr Tyr Phe Thr Gly Ile Trp Val Ile Ala His Glu Cys
 100 105 110
 Gly His Gln Ala Phe Ser Ala Ser Glu Ile Leu Asn Asp Thr Val Gly
 115 120 125
 Ile Ile Leu His Ser Leu Leu Phe Val Pro Tyr His Ser Trp Lys Ile
 130 135 140
 Thr His Arg Arg His His Ser Asn Thr Gly Ser Cys Glu Asn Asp Glu
 145 150 155 160
 Val Phe Thr Pro Thr Pro Arg Ser Val Val Glu Ala Lys His Asp His
 165 170 175
 Ser Leu Leu Glu Glu Ser Pro Leu Tyr Asn Leu Tyr Gly Ile Val Met
 180 185 190
 Met Leu Leu Val Gly Trp Met Pro Gly Tyr Leu Phe Phe Asn Ala Thr
 195 200 205
 Gly Pro Thr Lys Tyr Ala Gly Leu Ala Lys Ser His Phe Asn Pro Tyr
 210 215 220
 Ala Ala Phe Phe Leu Pro Lys Glu Arg Leu Ser Ile Trp Trp Ser Asp
 225 230 235 240
 Leu Cys Phe Leu Ala Ala Leu Tyr Gly Phe Gly Tyr Gly Val Ser Val
 245 250 255
 Phe Gly Leu Leu Asp Val Ala Arg His Tyr Ile Val Pro Tyr Leu Ile
 260 265 270
 Cys Asn Ala Tyr Leu Val Leu Ile Thr Tyr Leu Gln His Thr Asp Thr
 275 280 285
 Tyr Val Pro His Phe Arg Gly Asp Glu Trp Asn Trp Leu Arg Gly Ala
 290 295 300
 Leu Cys Thr Val Asp Arg Ser Phe Gly Ala Trp Ile Asp Ser Ala Ile
 305 310 315 320
 His His Ile Ala Asp Thr His Val Thr His His Ile Phe Ser Lys Thr
 325 330 335
 Pro Phe Tyr His Ala Ile Glu Ala Thr Asp Ala Ile Thr Pro Leu Leu
 340 345 350
 Gly Lys Tyr Tyr Leu Ile Asp Pro Thr Pro Ile Pro Leu Ala Leu Trp
 355 360 365
 Arg Ser Phe Thr His Cys Lys Tyr Val Glu Asp Asp Gly Asn Val Val
 370 375 380
 Phe Tyr Lys Arg Lys Leu Glu Glu Lys
 385 390

<210> 43

<211> 393

<212> PRT

<213> Saprolegnia diclina

<400> 43

Met Cys Lys Gly Gln Ala Pro Ser Lys Ala Asp Val Phe His Ala Ala
 1 5 10 15
 Gly Tyr Arg Pro Val Ala Gly Thr Pro Glu Pro Leu Pro Leu Glu Pro

20	25	30
Pro Thr Ile Thr Leu Lys Asp Leu Arg Ala Ala Ile Pro Ala His Cys		
35	40	45
Phe Glu Arg Ser Ala Ala Thr Ser Phe Tyr His Leu Ala Lys Asn Leu		
50	55	60
Ala Ile Cys Ala Gly Val Phe Ala Val Gly Leu Lys Leu Ala Ala Ala		
65	70	75
Asp Leu Pro Leu Ala Ala Lys Leu Val Ala Trp Pro Ile Tyr Trp Phe		
85	90	95
Val Gln Gly Thr Tyr Phe Thr Gly Ile Trp Val Ile Ala His Glu Cys		
100	105	110
Gly His Gln Ala Phe Ser Ala Ser Glu Ile Leu Asn Asp Thr Val Gly		
115	120	125
Ile Ile Leu His Ser Leu Leu Phe Val Pro Tyr His Ser Trp Lys Ile		
130	135	140
Thr His Arg Arg His His Ser Asn Thr Gly Ser Cys Glu Asn Asp Glu		
145	150	155
Val Phe Thr Pro Thr Pro Arg Ser Val Val Glu Ala Lys His Asp His		
165	170	175
Ser Leu Leu Glu Glu Ser Pro Leu Tyr Asn Leu Tyr Gly Ile Val Met		
180	185	190
Met Leu Leu Val Gly Trp Met Pro Gly Tyr Leu Phe Phe Asn Ala Thr		
195	200	205
Gly Pro Thr Lys Tyr Ala Gly Leu Ala Lys Ser His Phe Asn Pro Tyr		
210	215	220
Ala Ala Phe Phe Leu Pro Lys Glu Arg Leu Ser Ile Trp Trp Ser Asp		
225	230	235
Leu Cys Phe Leu Ala Ala Leu Tyr Gly Phe Gly Tyr Gly Val Ser Val		
245	250	255
Phe Gly Leu Leu Asp Val Ala Arg His Tyr Ile Val Pro Tyr Leu Ile		
260	265	270
Cys Asn Ala Tyr Leu Val Leu Ile Thr Tyr Leu Gln His Thr Asp Thr		
275	280	285
Tyr Val Pro His Phe Arg Gly Asp Glu Trp Asn Trp Leu Arg Gly Ala		
290	295	300
Leu Cys Thr Val Asp Arg Ser Phe Gly Ala Trp Ile Asp Ser Ala Ile		
305	310	315
His His Ile Ala Asp Thr His Val Thr His Ile Phe Ser Lys Thr		
325	330	335
Pro Phe Tyr His Ala Ile Glu Ala Thr Asp Ala Ile Thr Pro Leu Leu		
340	345	350
Gly Lys Tyr Tyr Leu Ile Asp Pro Thr Pro Ile Pro Leu Ala Leu Trp		
355	360	365
Arg Ser Phe Thr His Cys Lys Tyr Val Glu Asp Asp Gly Asn Val Val		
370	375	380
Phe Tyr Lys Arg Lys Leu Glu Glu Lys		
385	390	

<210> 44

<211> 359

<212> PRT

<213> Synechocystis sp.

<220>

<221> VARIANT

<222> (315)...(315)

<223> Xaa = Unknown or Other at position 315

<221> VARIANT

<222> (331)...(331)

<223> Xaa = Unknown or Other at position 331

<400> 44

```

Tyr Phe Phe Leu Asp Val Gly Leu Ile Ala Gly Phe Tyr Ala Leu Ala
 1           5           10           15
Ala Tyr Leu Asp Ser Trp Phe Phe Tyr Pro Ile Phe Trp Leu Ile Gln
          20           25           30
Gly Thr Leu Phe Trp Ser Leu Phe Val Val Gly His Asp Cys Gly His
          35           40           45
Gly Ser Phe Ser Lys Ser Lys Thr Leu Asn Asn Trp Ile Gly His Leu
          50           55           60
Ser His Thr Pro Ile Leu Val Pro Tyr His Gly Trp Arg Ile Ser His
65          70           75           80
Arg Thr His His Ala Asn Thr Gly Asn Ile Asp Thr Asp Glu Ser Trp
          85           90           95
Tyr Pro Val Ser Glu Gln Lys Tyr Asn Gln Met Ala Trp Tyr Glu Lys
          100          105          110
Leu Leu Arg Phe Tyr Leu Pro Leu Ile Ala Tyr Pro Ile Tyr Leu Phe
          115          120          125
Arg Arg Ser Pro Asn Arg Gln Gly Ser His Phe Met Pro Gly Ser Pro
          130          135          140
Leu Phe Arg Pro Gly Glu Lys Ala Ala Val Leu Thr Ser Thr Phe Ala
145          150          155          160
Leu Ala Ala Phe Val Gly Phe Leu Gly Phe Leu Thr Trp Gln Phe Gly
          165          170          175
Trp Leu Phe Leu Leu Lys Phe Tyr Val Ala Pro Tyr Leu Val Phe Val
          180          185          190
Val Trp Leu Asp Leu Val Thr Phe Leu His His Thr Glu Asp Asn Ile
          195          200          205
Pro Trp Tyr Arg Gly Asp Asp Trp Tyr Phe Leu Lys Gly Ala Leu Ser
          210          215          220
Thr Ile Asp Arg Asp Tyr Gly Phe Ile Asn Pro Ile His His Asp Ile
225          230          235          240
Gly Thr His Val Ala His His Ile Phe Ser Asn Met Pro His Tyr Lys
          245          250          255
Leu Arg Arg Ala Thr Glu Ala Ile Lys Pro Ile Leu Gly Glu Tyr Tyr
          260          265          270
Arg Tyr Ser Asp Glu Pro Ile Trp Gln Ala Phe Phe Lys Ser Tyr Trp
          275          280          285
Ala Cys His Phe Val Pro Asn Gln Gly Ser Gly Val Tyr Tyr Gln Ser
          290          295          300
Pro Ser Asn Gly Gly Tyr Gln Lys Lys Pro Xaa Leu Ile Leu Ile Glu
305          310          315          320
Ser Asn Gln His Arg Glu Gly Arg Gln Tyr Xaa Met Val Leu Leu Pro
          325          330          335
Ser Asp Arg Leu Met Arg Ser Met Glu Glu Val Lys Gln Ser His Ser
          340          345          350
Lys Arg Ser Ala Leu Asn Gln
          355

```

<210> 45

<211> 358

<212> PRT

<213> Saprolegnia diclina

<400> 45.

```

Met Thr Glu Asp Lys Thr Lys Val Glu Phe Pro Thr Leu Thr Glu Leu
 1          5          10          15
Lys His Ser Ile Pro Asn Ala Cys Phe Glu Ser Asn Leu Gly Leu Ser
 20          25          30
Leu Tyr Tyr Thr Ala Arg Ala Ile Phe Asn Ala Ser Ala Ser Ala Ala
 35          40          45
Leu Leu Tyr Ala Ala Arg Ser Thr Pro Phe Ile Ala Asp Asn Val Leu
 50          55          60
Leu His Ala Leu Val Cys Ala Thr Tyr Ile Tyr Val Gln Gly Val Ile
 65          70          75          80
Phe Trp Gly Phe Phe Thr Val Gly His Asp Cys Gly His Ser Ala Phe
 85          90          95
Ser Arg Tyr His Ser Val Asn Phe Ile Ile Gly Cys Ile Met His Ser
 100          105          110
Ala Ile Leu Thr Pro Phe Glu Ser Trp Arg Val Thr His Arg His His
 115          120          125
His Lys Asn Thr Gly Asn Ile Asp Lys Asp Glu Ile Phe Tyr Pro His
 130          135          140
Arg Ser Val Lys Asp Leu Gln Asp Val Arg Gln Trp Val Tyr Thr Leu
 145          150          155          160
Gly Gly Ala Trp Phe Val Tyr Leu Lys Val Gly Tyr Ala Pro Arg Thr
 165          170          175
Met Ser His Phe Asp Pro Trp Asp Pro Leu Leu Leu Arg Arg Ala Ser
 180          185          190
Ala Val Ile Val Ser Leu Gly Val Trp Ala Ala Phe Phe Ala Ala Tyr
 195          200          205
Ala Tyr Leu Thr Tyr Ser Leu Gly Phe Ala Val Met Gly Leu Tyr Tyr
 210          215          220
Tyr Ala Pro Leu Phe Val Phe Ala Ser Phe Leu Val Ile Thr Thr Phe
 225          230          235          240
Leu His His Asn Asp Glu Ala Thr Pro Trp Tyr Gly Asp Ser Glu Trp
 245          250          255
Thr Tyr Val Lys Gly Asn Leu Ser Ser Val Asp Arg Ser Tyr Gly Ala
 260          265          270
Phe Val Asp Asn Leu Ser His His Ile Gly Thr His Gln Val His His
 275          280          285
Leu Phe Pro Ile Ile Pro His Tyr Lys Leu Asn Glu Ala Thr Lys His
 290          295          300
Phe Ala Ala Ala Tyr Pro His Leu Val Arg Arg Asn Asp Glu Pro Ile
 305          310          315          320
Ile Thr Ala Phe Phe Lys Thr Ala His Leu Phe Val Asn Tyr Gly Ala
 325          330          335
Val Pro Glu Thr Ala Gln Ile Phe Thr Leu Lys Glu Ser Ala Ala Ala
 340          345          350
Ala Lys Ala Lys Ser Asp
 355

```

<210> 46

<211> 409

<212> PRT

<213> *Caenorhabditis elegans*

<220>

<221> VARIANT

<222> (389)...(389)

<223> Xaa = Unknown or Other at position 389

<400> 46

Val	Thr	Gly	Gly	Asp	Val	Leu	Val	Asp	Ala	Arg	Ala	Ser	Leu	Glu	Glu
1				5					10					15	
Lys	Glu	Ala	Pro	Arg	Asp	Val	Asn	Ala	Asn	Thr	Lys	Gln	Ala	Thr	Thr
			20					25					30		
Glu	Glu	Pro	Arg	Ile	Gln	Leu	Pro	Thr	Val	Asp	Ala	Phe	Arg	Arg	Ala
		35					40					45			
Ile	Pro	Ala	His	Cys	Phe	Glu	Arg	Asp	Leu	Val	Lys	Ser	Ile	Arg	Tyr
	50					55					60				
Leu	Val	Gln	Asp	Phe	Ala	Ala	Leu	Thr	Ile	Leu	Tyr	Phe	Ala	Leu	Pro
65					70					75					80
Ala	Phe	Glu	Tyr	Phe	Gly	Leu	Phe	Gly	Tyr	Leu	Val	Trp	Asn	Ile	Phe
				85				90						95	
Met	Gly	Val	Phe	Gly	Phe	Ala	Leu	Phe	Val	Val	Gly	His	Asp	Cys	Leu
			100					105					110		
His	Gly	Ser	Phe	Ser	Asp	Asn	Gln	Asn	Leu	Asn	Asp	Phe	Ile	Gly	His
		115				120						125			
Ile	Ala	Phe	Ser	Pro	Leu	Phe	Ser	Pro	Tyr	Phe	Pro	Trp	Gln	Lys	Ser
	130					135					140				
His	Lys	Leu	His	His	Ala	Phe	Thr	Asn	His	Ile	Asp	Lys	Asp	His	Gly
145					150					155					160
His	Val	Trp	Ile	Gln	Asp	Lys	Asp	Trp	Glu	Ala	Met	Pro	Ser	Trp	Lys
				165					170					175	
Arg	Trp	Phe	Asn	Pro	Ile	Pro	Phe	Ser	Gly	Trp	Leu	Lys	Trp	Phe	Pro
			180					185					190		
Val	Tyr	Thr	Leu	Phe	Gly	Phe	Cys	Asp	Gly	Ser	His	Phe	Trp	Pro	Tyr
		195					200					205			
Ser	Ser	Leu	Phe	Val	Arg	Asn	Ser	Asp	Arg	Val	Gln	Cys	Val	Ile	Ser
	210					215					220				
Gly	Ile	Cys	Cys	Cys	Val	Cys	Ala	Tyr	Ile	Ala	Leu	Thr	Ile	Ala	Gly
225					230					235					240
Ser	Tyr	Ser	Asn	Trp	Phe	Trp	Tyr	Tyr	Trp	Val	Pro	Leu	Ser	Phe	Phe
				245					250					255	
Gly	Leu	Met	Leu	Val	Ile	Val	Thr	Tyr	Leu	Gln	His	Val	Asp	Asp	Val
		260						265					270		
Ala	Glu	Val	Tyr	Glu	Ala	Asp	Glu	Trp	Ser	Phe	Val	Arg	Gly	Gln	Thr
		275					280					285			
Gln	Thr	Ile	Asp	Arg	Tyr	Tyr	Gly	Leu	Gly	Leu	Asp	Thr	Thr	Met	His
	290					295					300				
His	Ile	Thr	Asp	Gly	His	Val	Ala	His	His	Phe	Phe	Asn	Lys	Ile	Pro
305					310					315					320
His	Tyr	His	Leu	Ile	Glu	Ala	Thr	Glu	Gly	Val	Lys	Lys	Val	Leu	Glu
				325					330					335	
Pro	Leu	Ser	Asp	Thr	Gln	Tyr	Gly	Tyr	Lys	Ser	Gln	Val	Asn	Tyr	Asp
			340					345					350		
Phe	Phe	Ala	Arg	Phe	Leu	Trp	Phe	Asn	Tyr	Lys	Leu	Asp	Tyr	Leu	Val
		355					360					365			
His	Lys	Thr	Ala	Gly	Ile	Met	Gln	Phe	Arg	Thr	Thr	Leu	Glu	Glu	Lys
	370					375					380				
Ala	Lys	Ala	Lys	Xaa	Lys	Asn	Ile	Pro	Cys	Arg	Ser	Arg	Val	Gln	Gln
385					390					395					400
Gln	Leu	Leu	Arg	Phe	His	Arg	Phe	Cys							
				405											

<210> 47

<211> 333
 <212> PRT
 <213> *Saprolegnia diclina*

<400> 47

Met	Cys	Lys	Gly	Gln	Ala	Pro	Ser	Lys	Ala	Asp	Val	Phe	His	Ala	Ala
1				5					10					15	
Gly	Tyr	Arg	Pro	Val	Ala	Gly	Thr	Pro	Glu	Pro	Leu	Pro	Leu	Glu	Pro
			20					25					30		
Pro	Thr	Ile	Thr	Leu	Lys	Asp	Leu	Arg	Ala	Ala	Ile	Pro	Ala	His	Cys
		35					40					45			
Phe	Glu	Arg	Ser	Ala	Ala	Thr	Ser	Phe	Tyr	His	Leu	Ala	Lys	Asn	Leu
	50					55					60				
Ala	Ile	Cys	Ala	Gly	Val	Phe	Ala	Val	Gly	Leu	Lys	Leu	Ala	Ala	Ala
65					70					75					80
Asp	Leu	Pro	Leu	Ala	Ala	Lys	Leu	Val	Ala	Trp	Pro	Ile	Tyr	Trp	Phe
			85						90					95	
Val	Gln	Gly	Thr	Tyr	Phe	Thr	Gly	Ile	Trp	Val	Ile	Ala	His	Glu	Cys
			100					105					110		
Gly	His	Gln	Ala	Phe	Ser	Ala	Ser	Glu	Ile	Leu	Asn	Asp	Thr	Val	Gly
		115					120					125			
Ile	Ile	Leu	His	Ser	Leu	Leu	Phe	Val	Pro	Tyr	His	Ser	Trp	Lys	Ile
	130					135					140				
Thr	His	Arg	Arg	His	His	Ser	Asn	Thr	Gly	Ser	Cys	Glu	Asn	Asp	Glu
145					150					155					160
Val	Phe	Thr	Pro	Thr	Pro	Arg	Ser	Val	Val	Glu	Ala	Lys	His	Asp	His
				165					170					175	
Ser	Leu	Leu	Glu	Glu	Ser	Pro	Leu	Tyr	Asn	Leu	Tyr	Gly	Ile	Val	Met
			180					185					190		
Met	Leu	Leu	Val	Gly	Trp	Met	Pro	Gly	Tyr	Leu	Phe	Phe	Asn	Ala	Thr
		195					200					205			
Gly	Pro	Thr	Lys	Tyr	Ala	Gly	Leu	Ala	Lys	Ser	His	Phe	Asn	Pro	Tyr
	210					215					220				
Ala	Ala	Phe	Phe	Leu	Pro	Lys	Glu	Arg	Leu	Ser	Ile	Trp	Trp	Ser	Asp
225					230					235					240
Leu	Cys	Phe	Leu	Ala	Ala	Leu	Tyr	Gly	Phe	Gly	Tyr	Gly	Val	Ser	Val
				245					250					255	
Phe	Gly	Leu	Leu	Asp	Val	Ala	Arg	His	Tyr	Ile	Val	Pro	Tyr	Leu	Ile
			260					265					270		
Cys	Asn	Ala	Tyr	Leu	Val	Leu	Ile	Thr	Tyr	Leu	Gln	His	Thr	Asp	Thr
		275					280					285			
Thr	Pro	Leu	Leu	Gly	Lys	Tyr	Tyr	Leu	Ile	Asp	Pro	Thr	Pro	Ile	Pro
	290					295					300				
Leu	Ala	Leu	Trp	Arg	Ser	Phe	Thr	His	Cys	Lys	Tyr	Val	Glu	Asp	Asp
305					310					315					320
Gly	Asn	Val	Val	Phe	Tyr	Lys	Arg	Lys	Leu	Glu	Glu	Lys			
				325						330					

<210> 48
 <211> 412
 <212> PRT
 <213> *Gossypium hirsutum*

<220>

<221> VARIANT

<222> (9)...(9)

<223> Xaa = Unknown or Other at position 9

<221> VARIANT

<222> (403)...(403)

<223> Xaa = Unknown or Other at position 403

<400> 48

```

Leu Arg Val Ser Ser Thr Trp Arg Xaa Thr Ala Phe Phe Lys Ala Ser
1      5      10      15
Lys Met Gly Ala Gly Gly Arg Met Pro Ile Asp Gly Ile Lys Glu Glu
20      25      30
Asn Arg Gly Ser Val Asn Arg Val Pro Ile Glu Lys Pro Pro Phe Thr
35      40      45
Leu Gly Gln Ile Lys Gln Ala Ile Pro Pro His Cys Phe Arg Arg Ser
50      55      60
Leu Leu Arg Ser Phe Ser Tyr Val Val His Asp Leu Cys Leu Ala Ser
65      70      75      80
Phe Phe Tyr Tyr Ile Ala Thr Ser Tyr Phe His Phe Leu Pro Gln Pro
85      90      95
Phe Ser Tyr Ile Ala Trp Pro Val Tyr Trp Val Leu Gln Gly Cys Ile
100      105      110
Leu Thr Gly Val Trp Val Ile Ala His Glu Trp Gly His His Ala Phe
115      120      125
Arg Asp Tyr Gln Trp Val Asp Asp Thr Val Gly Leu Ile Leu His Ser
130      135      140
Ala Leu Leu Val Pro Tyr Phe Ser Trp Lys Ile Ser His Arg Arg His
145      150      155      160
His Ser Asn Thr Gly Ser Met Glu Arg Asp Glu Val Phe Val Pro Lys
165      170      175
Pro Lys Ser Lys Leu Ser Cys Phe Ala Lys Tyr Leu Asn Asn Pro Pro
180      185      190
Gly Arg Val Leu Ser Leu Val Val Thr Leu Thr Leu Gly Trp Pro Met
195      200      205
Tyr Leu Ala Phe Asn Val Ser Gly Arg Tyr Tyr Asp Arg Leu Ala Ser
210      215      220
His Tyr Asn Pro Tyr Gly Pro Ile Tyr Ser Asp Arg Glu Arg Leu Gln
225      230      235      240
Val Tyr Ile Ser Asp Thr Gly Ile Phe Ala Val Ile Tyr Val Leu Tyr
245      250      255
Lys Ile Ala Ala Thr Lys Gly Leu Ala Trp Leu Leu Cys Thr Tyr Gly
260      265      270
Val Pro Leu Leu Ile Val Asn Ala Phe Leu Val Leu Ile Thr Tyr Leu
275      280      285
Gln His Thr His Ser Ala Leu Pro His Tyr Asp Ser Ser Glu Trp Asp
290      295      300
Trp Leu Arg Gly Ala Leu Ser Thr Met Asp Arg Asp Phe Gly Val Leu
305      310      315      320
Asn Lys Val Phe His Asn Ile Thr Asp Thr His Val Ala His His Leu
325      330      335
Phe Ser Thr Met Pro His Tyr His Ala Met Glu Ala Thr Lys Ala Ile
340      345      350
Lys Pro Ile Leu Gly Lys Tyr Tyr Pro Phe Asp Gly Thr Pro Ile Tyr
355      360      365
Lys Ala Met Trp Arg Glu Ala Lys Glu Cys Leu Tyr Val Glu Pro Asp
370      375      380
Val Gly Gly Gly Gly Gly Gly Ser Lys Gly Val Phe Trp Tyr Arg Asn
385      390      395      400
Lys Phe Xaa Arg Pro Thr Asn Cys Leu Ile Ala Gly

```

24.

405

410

<210> 49
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Protein Motif 1 from Example 3

<400> 49
Thr Arg Ala Ala Ile Pro Lys His Cys Trp Val Lys
1 5 10

<210> 50
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Protein Motif 2 from Example 3

<400> 50
Ala Leu Phe Val Leu Gly His Asp Cys Gly His Gly Ser Phe Ser
1 5 10 15

<210> 51
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Protein Motif 3 from Example 3

<400> 51
Pro Tyr His Gly Trp Arg Ile Ser His Arg Thr His His Gln Asn
1 5 10 15

<210> 52
<211> 12
<212> PRT
<213> Artificial Sequence

<220>
<223> Protein Motif 4 from Example 3

<221> VARIANT
<222> (5)...(5)
<223> Xaa = D or H at position 5

<221> VARIANT
<222> (7)...(7)
<223> Xaa = D or Y at position 7

<400> 52
Gly Ser His Phe Xaa Pro Xaa Ser Asp Leu Phe Val
1 5 10

<210> 53
<211> 13
<212> PRT
<213> Artificial Sequence

<220>
<223> Protein Motif 5 from Example 3

<221> VARIANT
<222> (3)...(3)
<223> Xaa = Y or F at position 3

<221> VARIANT
<222> (4)...(4)
<223> Xaa = L or V at position 4

<221> VARIANT
<222> (11)...(11)
<223> Xaa = L or I at position 11

<400> 53
Trp Ser Xaa Xaa Arg Gly Gly Leu Thr Thr Xaa Asp Arg
1 5 10

<210> 54
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Protein Motif 6 from Example 3

<400> 54
His His Asp Ile Gly Thr His Val Ile His His Leu Phe Pro Gln
1 5 10 15

<210> 55
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Protein Motif 7 from Example 3

<221> VARIANT
<222> (2)...(2)
<223> Xaa = L or F at position 2

<221> VARIANT
<222> (5)...(5)
<223> Xaa = Q or K at position 5

<221> VARIANT
<222> (12)...(12)
<223> Xaa = V or I at position 12

26.

<400> 55

His Xaa Phe Pro Xaa Ile Pro His Tyr His Leu Xaa Glu Ala Thr
1 5 10 15

<210> 56

<211> 15

<212> PRT

<213> Artificial Sequence

<220>

<223> Protein Motif 8 from Example 3

<221> VARIANT

<222> (3)...(3)

<223> Xaa = A or I at position 3

<221> VARIANT

<222> (6)...(6)

<223> Xaa = L or F at position 6

<400> 56

His Val Xaa His His Xaa Phe Pro Gln Ile Pro His Tyr His Leu
1 5 10 15

<210> 57

<211> 17

<212> PRT

<213> Artificial Sequence

<220>

<223> Protein Motif 1 from Example 7

<221> VARIANT

<222> (2)...(2)

<223> Xaa = N or E at position 2

<221> VARIANT

<222> (10)...(10)

<223> Xaa = D or E at position 10

<221> VARIANT

<222> (11)...(11)

<223> Xaa = A or C at position 11

<400> 57

Pro Xaa Phe Thr Ile Lys Glu Ile Arg Xaa Xaa Ile Pro Ala His Cys
1 5 10 15
Phe

<210> 58

<211> 16

<212> PRT

<213> Artificial Sequence

<220>

<223> Protein Motif 2 from Example 7

<221> VARIANT
<222> (3)...(3)
<223> Xaa = H or F at position 3

<221> VARIANT
<222> (11)...(11)
<223> Xaa = V or Y at position 11

<221> VARIANT
<222> (13)...(13)
<223> Xaa = I or L at position 13

<221> VARIANT
<222> (16)...(16)
<223> Xaa = A or L at position 16

<400> 58
Met Pro Xaa Tyr His Ala Glu Glu Ala Thr Xaa His Xaa Lys Lys Xaa
1 5 10 15

<210> 59
<211> 15
<212> PRT
<213> Artificial Sequence

<220>
<223> Protein Motif 3 from Example 7

<221> VARIANT
<222> (2)...(2)
<223> Xaa = L or V at position 2

<221> VARIANT
<222> (5)...(5)
<223> Xaa = A or I at position 5

<221> VARIANT
<222> (6)...(6)
<223> Xaa = C or M or A at position 6

<221> VARIANT
<222> (9)...(9)
<223> Xaa = V or I at position 9

<221> VARIANT
<222> (11)...(11)
<223> Xaa = L or G or C at position 11

<400> 59
Pro Xaa Tyr Trp Xaa Xaa Gln Gly Xaa Val Xaa Thr Gly Val Trp
1 5 10 15

<210> 60
<211> 15
<212> PRT
<213> Artificial Sequence

<220>

<223> Protein Motif 4 from Example 7

<221> VARIANT

<222> (6)...(6)

<223> Xaa = L or F at position 6

<221> VARIANT

<222> (9)...(9)

<223> Xaa = T or Q at position 9

<400> 60

His	Val	Ala	His	His	Xaa	Phe	Ser	Xaa	Met	Pro	His	Tyr	His	Ala
1				5					10					15

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
7 August 2003 (07.08.2003)

PCT

(10) International Publication Number
WO 03/064596 A3

(51) International Patent Classification⁷:
C12N 15/82, 15/74, 5/04, A01H 5/00

C07H 21/02,

(74) Agents: BECKER, Cheryl, L. et al.; D-377 AP6A-1, 100
Abbott Park Road, Abbott Park, IL 60064-6008 (US).

(21) International Application Number: PCT/US03/01698

(81) Designated States (*national*): CA, JP, MX, NO.

(22) International Filing Date: 21 January 2003 (21.01.2003)

(84) Designated States (*regional*): European patent (AT, BE,
BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT,
LU, MC, NL, PT, SE, SK, TR).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
10/060,793 30 January 2002 (30.01.2002) US

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

(71) Applicant: ABBOTT LABORATORIES [US/US];
D-377 AP6A-1, 100 Abbott Park Road, Abbott Park, IL
60064-6008 (US).

(88) Date of publication of the international search report:
4 December 2003

(72) Inventors: MUKERJI, Pradip; 1069 Arcaro Drive,
Columbus, OH 43230 (US). PEREIRA, Suzette, L.; 710
Westray Drive, Westerville, OH 43081 (US). HUANG,
Yung-Sheng; 2462 Danvers Court, Upper Arlington, OH
43220 (US).

*For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.*

WO 03/064596 A3

(54) Title: DESATURASE GENES, ENZYMES ENCODED THEREBY, AND USES THEREOF

(57) Abstract: Disclosed are isolated polynucleotides encoding an omega-3 desaturase and a delta-12 desaturase, the enzymes encoded by the isolated polynucleotides, vectors containing the isolated polynucleotides, transgenic hosts that contain the isolated polynucleotides that express the enzymes encoded thereby, methods for producing the desaturase enzymes, and method of using the enzymes to make polyunsaturated fatty acids. The isolated polynucleotides are derived from a fungus, *Saprolegnia diclina* (ATCC 56851). In particular, omega-3-desaturase may be utilized, for example, in the conversion of arachidonic acid (AA) to eicosapentaenoic acid (EPA). Delta-12 desaturase may be used, for example, in the conversion of oleic acid (OA) to linoleic (LA). EPA or polyunsaturated fatty acids produced therefrom may be added to pharmaceutical compositions, nutritional compositions, animal feeds, as well as other products such as cosmetics.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/01698

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C07H 21/02; C12N 15/82, 15/74, 5/04; A01H 5/00

US CL : 536/23.2; 435/468, 471, 419, 252.3; 800/281, 298

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 536/23.2; 435/468, 471, 419, 252.3; 800/281, 298

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
SEQUENCE, WEST, DIALOG

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,658,767 A (KYLE et al) 19 August 1997 (19.08.1997), columns 1-4.	1-31
A	US 6,150,144 A (AKIMOTO et al) 21 November 2000 (21.11.2000), columns 1-6.	1-31

☐ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier application or patent published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

12 September 2003 (12.09.2003)

Date of mailing of the international search report

14 OCT 2003

Name and mailing address of the ISA/US

Mail Stop PCT, Attn: ISA/US
Commissioner for Patents
P.O. Box 1450
Alexandria, Virginia 22313-1450

Facsimile No. (703)305-3230

Authorized officer

Elizabeth F. McElwain

Telephone No. 703-308-0196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US03/01698

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claim Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claim Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claim Nos.: 32-39
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
Please See Continuation Sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐
☒

The additional search fees were accompanied by the applicant's protest.
No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

PCT/US03/01698

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

This application contains the following inventions or groups of inventions which are not so linked as to form a single general inventive concept under PCT Rule 13.1. In order for all inventions to be examined, the appropriate additional examination fees must be paid.

Group I, claim(s) 1-5, 11-18 and 20-31, to the extent the claims are drawn to the omega-3 desaturase of SEQ ID NO: 25 encoding SEQ ID NO: 26.

Group II, claim(s) 1-5, 11-18 and 20-31, drawn to the delta-12 desaturase of SEQ ID NO: 41 encoding SEQ ID NO: 42.

Group III, claim(s) 6, 7, 8 and 10, drawn to the polypeptide of SEQ ID NO: 26.

Group IV, claim(s) 6 and 9, drawn to the polypeptide of SEQ ID NO: 42.

Group V, claim(s) 19, drawn to one or more plant oils.

The inventions listed as Groups I-V do not relate to a single general inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: the inventions of Groups I and II are drawn to DNA coding for enzymatically distinct polypeptides, which differ structurally and functionally. Groups III and IV are each drawn to distinct polypeptides that differ structurally and functionally from one another and from the DNA of Groups I and II. In addition, the oil of Group V differs chemically, structurally and functionally from the products of Groups I-IV and there is no corresponding special technical feature shared by these Groups.

THIS PAGE BLANK (USPTO)